

ACCEPTED
802-26

[802-26]

Miller's
ROTENONE 100

GENERAL DIRECTIONS

1. Dust lightly on first appearance of insect or insect damage.
2. Repeat at 4 to 7-day intervals until under control.
3. Frequent light dustings are more effective than single heavy dustings.
4. Do not apply to food crops within one day of harvest.

VEGETABLES

ASPARAGUS—Asparagus Beetle
CUCUMBER—Fly Beetles, Mexican Bean Beetles, Cucumber Beetles
CABBAGE, **CARLEFLOWER**, **BROCCOLI**, **BRUSSELS SPROUTS**, **COLLARD**, **KOHLEBI**, **KALE**—Fly, Cabbage Worms and Loopers, Vegetable Weevil
MELON, **SQUASH**—Cucumber Beetles, Fleas
CABBAGE—Cabbage Looper
SPINACH—Aphids, Flea Weevil
 For Flea Weevil control, dust on foliage during early bloom period before eggs are laid.
TOMATOES, **PLANTAIN**, **PEPPERS**—Aphids, Colorado Potato Beetles, Flea Beetles
PEPPERS—Aphids, Flea Beetles, Vegetable Weevil

FLOWERS

ROSE—Aphids, Flea Beetles, Cucumber Beetles, Flea Weevil, Green and Rose

NET WT.
1 LB.



Rotenone
100 (1% Rotenone Dust)
Insecticide



CAUTION Keep out of the reach of children
See right side panel for additional cautions.
E.P.A. Reg. No. 802-26-AA

Miller's
ROTECNE 100

SMALL FRUITS

BLACKBERRIES, LOGANBERRIES, RASPBERRIES

1. For Raspberry Cane Borer and Rednecked Cane Borer, dust foliage just before plants bloom. Repeat in about 2 weeks.
2. For Raspberry Fruitworm, dust foliage 7 days after first blossom. Repeat in 10 days and again 10 days later.

STRAWBERRIES—Spittle Bugs, Flea Beetles.

CAUTION

Harmful if swallowed. Avoid breathing dust. Avoid contact with skin; wash after handling.
 This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.
 Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes.

Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with chemicals not specifically recommended and other important factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given hereon.

THE CHAS. H. LILLY CO.
PORTLAND, OREGON
EPA Est. 802-OR-1

RECEIVED
17 OCT 1974
FEDERAL BUREAU OF INVESTIGATION
WASHINGTON, DC

ACCEPTED
802-26

[802-26]

NET WT.
1 LB.

Miller's ROTENONE 100

GENERAL DIRECTIONS

1. Dust lightly on first appearance of insect or insect damage.
2. Repeat at 4 to 7-day intervals until under control.
3. Frequent light dustings are more effective than single heavy dustings.
4. Do not apply to food crops within one day of harvest.

VEGETABLES

- ASPARAGUS - Asparagus Beetle
- BEANS, BEETS - Flea Beetles, Mexican Bean Beetles, Cucumber Beetles.
- CABBAGE, CAULIFLOWER, BROCCOLI, BRUSSELS SPROUTS, COLLARDS, KOHLRABI, KALE - Flea Beetles, Cabbage Worms and Loopers, Vegetable Weevil.
- CARROTS - Vegetable Weevil
- CUCUMBERS, MELONS, SQUASH - Cucumber Beetles, Pickleworms.
- LETTUCE - Cabbage Looper.
- PEAS - Aphids, Pea Weevil.
For Pea Weevil control dust on foliage during early bloom period before eggs are laid.
- POTATOES, TOMATOES, EGGPLANTS, PEPPERS - Aphids, Colorado Potato Beetles, Flea Beetles.
- RUTABAGAS - Aphids, Flea Beetles, Vegetable Weevil.

FLOWERS

- FLOWERS such as Bees, Chrysanthemums, Dahlias, Peonies, Sweet Peas, Phlox, Delphiniums, Mari-golds, Zinnias - Flea Beetles, Cucumber Beetles (Diabrotica), Aphids (Green and Rosy).



Rotenone 100 (1% Rotenone) Insect



FOR FLOWERS, VEGETABLES
AND SMALL FRUITS

CAUTION Keep out of the reach of children.
See right side panel for additional information.
E.P.A. Reg. No. 802-26

Miller's ROTENONE 100

SMALL FRUITS

BLACKBERRIES, LOGANBERRIES, RASPBERRIES

1. For Raspberry Cane Borer and Rednecked Cane Borer, dust foliage just before plants bloom. Repeat in about 2 weeks.
2. For Raspberry Fruitworm, dust foliage 7 days after first blossom. Repeat in 10 days and again 10 days later.

STRAWBERRIES—Spittle Bugs, Flea Beetles.

CAUTION

Harmful if swallowed. Avoid breathing dust. Avoid contact with skin; wash after handling.

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes.

NOTICE: Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

MP-10J