

STORAGE AND DISPOSAL

Store this product in its original container and keep in a locked storage area out of reach of children and domestic animals. Do not reuse container. Securely wrap container in several layers of newspaper and discard in trash.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Avoid breathing dust. Avoid contact with skin; wash after handling.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Filled by weight not by volume. Product may settle after packaging.

NOTICE: Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

THE CHAS. H. LILLY CO.
PORTLAND, OREGON 97218

EPA Reg No 802-26

75152-1

EPA Est 802-OR-1

Label Received 10/4/88
NOT reviewed per Practice 88-6

LILLY/MILLER.

Rotenone 100 Insecticide (1% Rotenone Dust)



ACTIVE INGREDIENTS:	
Rotenone	1.00%
Other Cube Resins	1.00%
INERT INGREDIENTS 98.00%	

For Flowers, Vegetables, Small Fruits

Keep out of reach of children

CAUTION

See right panel for additional precautionary statements

NET WEIGHT 1 LB.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

1. Dust lightly on first appearance of insect or insect damage.
2. Repeat at 4 to 7 day intervals until under control.
3. Frequent light dustings are more effective than single heavy dustings.
4. Do not apply to food crops within one day of harvest.

VEGETABLES

ASPARAGUS—Asparagus Beetle.

BEANS, BEETS—Flea Beetles, Mexican Bean Beetles, Cucumber Beetles.

CABBAGE, CAULIFLOWER, BROCCOLI, BRUSSELS SPROUTS, COLLARDS, KOHLRABI, KALE—Flea Beetles, Imported Cabbageworms, Cabbage Loopers, Vegetable Weevil.

CARROTS—Vegetable Weevil.

CUCUMBERS, MELONS, SQUASH—Cucumber Beetles, Pickleworms.

LETUCE—Cabbage Looper.

PEAS—Aphids, Pea Weevil. For Pea Weevil control dust on foliage during early bloom period before eggs are laid.

POTATOES, TOMATOES, EGGPLANTS, PEPPERS—Aphids, Colorado Potato Beetles, Flea Beetles.

RUTABAGAS—Aphids, Flea Beetles, Vegetable Weevils.

FLOWERS

Flowers such as Roses, Chrysanthemums, Dahlias, Peonies, Sweet Peas, Phlox, Delphiniums, Marigolds, Zinnias—Flea Beetles, Cucumber Beetles (Diabrotica), Aphids (Green and Rosy).

SMALL FRUITS

BLACKBERRIES, LOGANBERRIES, RASPBERRIES

For Raspberry Cane Borer and Rednecked Cane Borer, dust foliage just before plants bloom. Repeat in about 2 weeks. For Raspberry Fruitworm, dust foliage 7 days after first blossom. Repeat in 10 days and again 10 days later.

STRAWBERRIES—Spittlebugs, Flea Beetles.

STORAGE AND DISPOSAL

Store this product in its original container and keep in a locked storage area out of reach of children and domestic animals. Do not reuse container. Securely wrap container in several layers of newspaper and discard in trash.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Avoid breathing dust. Avoid contact with skin; wash after handling.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Filled by weight not by volume. Product may settle after packaging.

NOTICE: Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

THE CHAS. H. LILLY CO.
PORTLAND, OREGON 97208

EPA Reg. No. 802-26

75152-1

EPA Est. 802-OR 1


LILLY/MILLER.

Rotenone 100 Insecticide (1% Rotenone Dust)



ACTIVE INGREDIENTS:

Rotenone	1.00%
Other Cube Resins	1.00%
INERT INGREDIENTS	98.00%

For Flowers, Vegetables, Small Fruits

Keep out of reach of children

CAUTION

See right panel for additional precautionary statements

NET WEIGHT 1 LB.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

1. Dust lightly on first appearance of insect or insect damage.
2. Repeat at 4 to 7 day intervals until under control.
3. Frequent light dustings are more effective than single heavy dustings.
4. Do not apply to food crops within one day of harvest.

VEGETABLES

ASPARAGUS—Asparagus Beetle.

BEANS, BEETS—Flea Beetles, Mexican Bean Beetles, Cucumber Beetles.

CABBAGE, CAULIFLOWER, BROCCOLI, BRUSSELS SPROUTS, COLLARDS, KOHLRABI, KALE—Flea Beetles, Imported Cabbageworms, Cabbage Loopers, Vegetable Weevil.

CARROTS—Vegetable Weevil.

CUCUMBERS, MELONS, SQUASH—Cucumber Beetles, Pickleworms.

LETTUCE—Cabbage Lopper.

PEAS—Aphids, Pea Weevil. For Pea Weevil control dust on foliage during early bloom period before eggs are laid.

POTATOES, TOMATOES, EGGPLANTS, PEPPERS—Aphids, Colorado Potato Beetles, Flea Beetles.

RUTABAGAS—Aphids, Flea Beetles, Vegetable Weevils.

FLOWERS

Flowers such as Roses, Chrysanthemums, Dahlias, Peonies, Sweet Peas, Phlox, Delphiniums, Marigolds, Zinnias—Flea Beetles, Cucumber Beetles (Diabrotica), Aphids (Green and Rosy).

SMALL FRUITS

BLACKBERRIES, LOGANBERRIES, RASPBERRIES

For Raspberry Cane Borer and Rednecked Cane Borer, dust foliage just before plants bloom. Repeat in about 2 weeks. For Raspberry Fruitworm, dust foliage 7 days after first blossom. Repeat in 10 days and again 10 days later.

STRAWBERRIES—Spittlebugs, Flea Beetles.