



# CRATER 98

DUSTING SULPHUR

JAN 31 1974

Active ingredient, Sulphur. . . . .	98%
Inert ingredients . . . . .	2%

### CAUTION — Keep out of reach of children

Sulphur will cause severe injury to sulphur-sensitive crops. Do not use on Apricots, Anjou and Comice Pears, Cranberries, Cucumbers, Squash and certain varieties of Melons, Filberts, Spinach and other sensitive plants.

Do not apply when shade temperatures exceed, or are likely to exceed, 90°F. Do not use sulphur within 30 days of an oil spray. Do not reuse container. Destroy when empty.

If crop to be treated will be used for canning or freezing purposes, consult food processor before applying sulphur.

### DIRECTIONS FOR USE

CHERRIES, PEACHES, PLUMS, PRUNES --- Brown Rot Blossom Blight  
Use 50 lbs. per acre. Apply at Popcorn, Full Bloom, and Petal Fall stages of blossom development.

CHERRIES, PEACHES, PLUMS, PRUNES - Brown Rot in Fruit  
Use 50 lbs. per acre. Make first application about 5 weeks before harvest or earlier if the disease appears. Repeat at weekly intervals with last application just before harvest.

GRAPES - Powdery Mildew, Rust  
Use 20 to 35 lbs. per acre. Apply when new shoots are 6 to 10 in. long; just before blossoming; when berries are set but no more than 1/4 grown; when berries are 1/2 to full

grown; just as grapes begin to ripen; and repeat as necessary for control.

ALFALFA, CLOVER - Spider Mites, Powdery Mildew  
Use 15 to 25 lbs. per acre. Apply when mites appear and before or when mildew appears. Repeat as necessary.

BEANS - Powdery Mildew, Spider Mites, Rust  
Use 15 to 30 lbs. per acre. Apply when mites appear and before or when mildew appears. Repeat as necessary for control. For Rust, make first application as soon as first spots appear. Continue dusting at 7 day intervals until blossom time.

GRASS SEED CROPS - Timothy Mites  
Use 30 lbs. per acre. Apply when mites appear and repeat as necessary for control.

PEAS - Powdery Mildew  
Use 15 to 20 lbs. per acre. Apply before or when disease appears and repeat as necessary for control.

ORNAMENTALS - Make full coverage application on both sides of leaves. Keep new growth protected. Some burning may occur when weather is warm.

Chrysanthemums - Powdery Mildew, Rust  
Do not apply when mums are flowering.  
Hollyhock - Rust  
Roses, Perennials, Shrubs - Powdery Mildew, Rust

NOTICE: Follow directions carefully. The use and method of application, weather and temperature may vary with the chemicals not specifically recommended. All other applications, uses, and methods of this product are beyond the control of the seller. Buyer assumes all risks of use. Storage and handling instructions are not to be taken as an acceptance with respect to the product.

E.P.A. Reg. No. 802-41-AA

MP-5H

Manufactured for ...

THE CHAS. H. LILLY CO.  
PORTLAND, OREGON

NET WT. 50 LBS.

