

[802-44]

# Miller's ROTEFIVE

**GENERAL DIRECTIONS**—do not apply to food crops within one day of harvest.

**MIXING AND SPRAYING DIRECTIONS**

1. Use 3 level tablespoons per gallon of water (3 lbs. per 100 gallons).
2. Make a thin paste with small amount of water and then add balance of water required.
3. Apply a minimum of 200 gallons of spray per acre every 4 to 7 days until insects are under control. Repeat after rains.
4. For Vegetables and Flowers, apply as per directions given above, for the control of insects listed.
5. For Fruits and Berries, use the above amounts for spraying and apply as per specific directions given for each crop.

### VEGETABLES

**SPINACH**—Aparagus Beetles.  
**BEANS**—Aparagus Beetles, Mexican Bean Beetles, Cucumber Beetles.  
**CABBAGE, CAULIFLOWER, BROCCOLI, BRUSSELS SPROUTS, COLLARDS, KOUZARI, KALE**—Aparagus Beetles, Imported Cabbage Worm, Cabbage Looper, Vegetable Weevils.  
**CARROTS**—Vegetable Weevils.  
**CELERY**—Aparagus.  
**CHAMBERS, MELONS, SQUASH**—Cucumber Beetles, Pickleworms.  
**LETTUCE**—Cabbage Looper.  
**PEAS**—Aparagus, Pea Weevil.  
 For Apples, follow directions above. For Pea Weevil, spray on foliage during early bloom period before eggs are laid.  
**POTATOES, TOMATOES, EGGPLANTS, PEPPERS**—Aparagus, Colorado Potato Beetles, Flea Beetles.  
**KUJABAS, TURNIPS**—Aparagus, Flea Beetles, Vegetable Weevils.

### FLOWERS

**ROSE, ANEMONE, HYDRANGEA, CHERRY, PEONIES, SYRINGA, BEARS, NIGELLA, DEUTERIUMS**  
 Colorado Potato Beetles, Flea Beetles, Imported Cabbage Worm, Green and Red Leafhoppers.


### FRUITS AND BERRIES

**CHERRIES**—Cherry Fruit Fly.  
 Make first application when fly first emerges (consult State Agricultural Extension Service Authorities for exact timing). Spray to cover entire tree. Repeat applications at 7 to 10 day intervals and after rains.  
**BLACKBERRIES, BOYBERRIES, DEWBERRIES, LOGANBERRIES, RASPBERRIES**—Raspberry Fruitworm, Red-necked Cane Borer.  
 Make first application for Fruitworm control when blossom buds first appear and repeat just before blossoms open. For Red-necked Cane Borer, make first application as above and second application about 2 weeks later. Spray to thoroughly cover all of the bush and both sides of the foliage. Repeat applications after rains.  
**CURRENTS, GOOSEBERRIES**—Gooseberry Fruitworms, Imported Current Worm.  
 Make first application when Fruitworm webbing is observed and 7 to 10 days later. For Current worm, make application when worms appear. Repeat as needed and after rains.  
**STRAWBERRIES**—Spittle Bug, Flea Beetles.  
 Make first application when insects are first seen. Repeat as needed and after rains.

### CAUTION

Harmful if swallowed or absorbed through skin. Avoid breathing dust. Avoid contact with skin. Wash after handling. Wear clean clothing.  
 This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated.  
 Rinse equipment and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes, or by burning (keep out of smoke). Do not contaminate water by cleaning of equipment or disposal of wastes.  
 Apply this product only as specified on this label.

**NOTICE:** Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of E.I. du Pont. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.



# ROTEFIVE

(5% Rotenone Spray Powder)

## INSECTICIDE

for  
**VEGETABLES**  
**SMALL FRUITS • FLOWERS**

Active ingredients:

Rotenone .....	5%
Other Cube Resins .....	5%
Inert ingredients .....	90%
Total.....100%	

### CAUTION — Keep out of reach of children

See back panel for additional cautions

E.P.A. Reg. No. 802-44-AA

## NET WEIGHT \_\_\_\_\_ LBS.

THE CHAS. H. LILLY CO.  
 PORTLAND, OREGON  
 E.P.A. Est. 802-OR-1

Miller's