802-44

LILLY MILLER ROTEFIVE (5% Rotenone Spray Powder)

ACTIVE INGREDIENTS: Rotenone....5% Other Cube Resins....5% INERT INGREDIENTS...90% Total....100%

INSECTICIDE for VEGETABLES, SMALL FRUITS, FLOWERS

Keep out of reach of children CAUTION See back panel for additional precautionary statements.

NET WEIGHT 4 LBS.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION: Harmful if swallowed. Avoid breathing dust. Avoid contact with skin, eyes or clothing. In case of contact, immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists.

Received by Votification

pr. P.R. Notice 88-6 11/4/88

802-44 107-3

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product through any type of irrigation system.

RE-ENTRY STATEMENT

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons. Do not enter treated areas without protective clothing until spray has dried. Because certain states. May require more restrictive re-entry intervals for various crops treated with this product, consult your State Department of Agriculture Bor. further information. Written or oral warnings must be given to workers who are expected to be in a treated area or an area about to be treated. Oral warnings must be given if written warnings cannot be mederstood by workers and must be in a language they understand. Give specific owal warnings which inform workers of areas or fields that may, be be entered without specific protective clothing, the period of time field must be vacated, and appropriate actions to take in case of accidental explosure. Written warnings must include the following information: CAUTION. Frea treated with ROTEFIVE on (date and time of application). Do not on the specific or other specific or

2053

without full protective clothing until sprays have dried.

Do not apply to food crops within one day of harvest.

MIXING AND SPRAYING DIRECTIONS

 Use 3 level tablespoons per gallon of water (3 lbs. per 100 gallons)
Make a thin paste with small amount of water and then add balance of water required.
Apply a minimum of 200 gallons of spray per acre every 4 to 7 days until insects are under control. Repeat after rains.
For Vegetables and Flowers, apply as per directions given above, for the control of insects listed.
For Fruits and Berries, use the above amounts for spraying and apply as per specific directions given for each crop.

VEGETABLES

ASPARAGUS-Asparagus Beetles. BEANS, BEETS-Flea Beetles, Mcmican Bean Beetles, Cucumber Beetles. CABBAGE, CAULIFLOWER, BROCCOLI, BRUSSELS SPROUTS, COLLARDS, KOHLRABI, KALE-Flea Beetles, Imported Cabbageworms, Cabbage Loopers, Vegetable Weevils. CARROTS-Vegetable Weevils. CELERY- Aphids. CUCUMBER, MELONS, SQUASH-Cucumber Beetles, Pickleworms. LETTUCE-Cabbage Loopers. PEAS-Aphids, Pea Weevil. For Aphids, follow directions above. For Pea Weevil, spray on foliage during early bloom period before eggs are laid. POTATOES, TOMATOES, EGGPLANTS, PEPPERS-Aphids, Colorado Potato Beetles, Flea Beetles. RUTABAGAS, TURNIPS-Aphids, Flea Beetles, Vegetable Weevils.

FLOWERS

ROSES, CHRYSANTHEMUMS, DAHLIAS, PEONIES, SWEET PEAS, PHLOX, DELPHINIUMS, MARIGOLDS, ZINNIAS-Flea Beetles, Cucumber Beetles, Aphids (Green and Rosy).

FRUITS AND BERRIES CHERRIES- Cherry Fruit Fly. Make first application when fly first emerges (consult State Agricultural Extension Service Authorities for exact timing). Spray to cover entire tree. Repeat applications at 7 to 10 day intervals and after rains.

BLACKBERRIES, BOYSENBERRIES, DEWBERRIES, LOGANBERRIES, RASPBERRIES-Raspberry Fruitworm, Red-necked Cane Borer. Make first application for Fruitworm control when blossom buds first appear and repeat just before blossoms open. For Red-necked Cane Borer, make first application as above and second application about 2 weeks later. Spray to thorbughly cover all of the bush and both sides of the foliage. Repeat applications after rains.

CURRANTS, GOOSEBERRIES-GOOSEberry Fruitworms, Imported Currant Worm. Make first application when Fruitworm webbing is observed and 7 to 10 days later. For Currant Worm, make application when worms appear, Repeat as needed and after rains.

STRAWBERRIES-Spittle Bugs, Flea Beetles. Make first application when insects are first seen. Repeat as needed and after rains.

:..:

••••

....

....

.

.....

STORAGE AND DISPOSAL

STORAGE: Store in a cool, dry place. DISPOSAL: Do not contaminate water, food or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

LIMIT OF WARRANTY AND LIABILITY

The Chas. *i.* LILLY Co. (LILLY) warrants that this product conforms to the chemical or other description on the label (and/or instructions) and is reasonably fit for the purposes set forth on the label (and/or instructions) when used strictly in accordance with the label (and/or instructions) under the conditions described there. NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS FOR A PARTICULAR PURPOSE OR MERCHANTABILITY IS MADE. This warranty is also subject to the following conditions: *Buyer and all users shall promptly notify LILLY of any claims whether based in contract, negligence, strict liability, other tort, or otherwise.

*THE EXCLUSIVE REMEDY OF THE USER OR BUYER AND THE LIMIT OF LIABILITY OF LILLY FOR ANY LOSS, INJURY OR DAMAGE RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT (including claims based upon ntract, negligence, strict liability, other tort, or otherwise) SHALL BE:

1. The purchase price paid for the product; or at the election of LILLY

2. The replacement of such quantity of the product as is involved. IN NO EVENT SHALL LILLY BE LIABLE FOR ANY INCIDENTAL OR CONSEQUENTIAL DAMAGES.

*LILLY agents have no authority to give warranties or make representations on its behalf, concerning the use, quality or performance of this product, or in any other respect. *The use, handling, storage, and other influencing factors are beyond

the control of LILLY. Buyer assumes all risk of loss, injury, or damage due to use, handling, storage and other influencing factors, including but not limited to, incompatability with other products, and weather conditions.

NOTICE: Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buver assumes and risks of use, storage or handling of this material not in strict accordance with directions given herewith.

Filled by weight not by volume. Froduct may settle after packaging.

THE CHAS. H. LILLY CO. PORTLAND, OREGON 97218 EPA Reg. No. 802-44 EPA Est. 802-0R-1 75104