

SUGGESTED USES

GENERAL—Use the amount given per acre in 6 to 10 gallons of water by air (20 to 50 gals. by ground equipment) to provide thorough coverage.

WIREWORMS, WHITE GRUBS—Beans, Beets, Blackberries, Blueberries, Broccoli, Cabbage, Cauliflower, Celery, Corn, Loganberries, Onions, Peppers, Potatoes, Strawberries, Sugar Beets, Turnips.

Use 2½ to 5 pints per acre. Apply to the soil surface and work into top 8 or 9 inches before planting.

FLEA BEETLES—Beans, Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Kohlrabi

Use 1 to 2 pints per acre. Repeat applications if necessary.

ONION THRIPS—Onions

Use 1 to 2 pints per acre as a foliage spray. Repeat applications if necessary. Do not apply to green or spring onions.

ROOT MAGGOTS—Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Kohlrabi, Radishes, Rutabagas, Turnips

Use 1 to 2 pints per acre worked into top 3 or 4 inches of soil, as a pre-planting, planting time or transplanting treatment.

ANTS—Lawns, Ornamental Turfs, Parks, Cemeteries, Nursery Stock, Ornamentals.

Use 1 pint per acre. Apply directly on ants, along their trails and douse hills or nests.

Birds feeding on treated areas may be harmed.

Keep children and pets off treated areas until this material is washed into the soil and grass is dry.

TERMITES (Subterranean)

Use 4¼ qts. Miller's Chlordane 8E in 98-15/16 gals. of water. This makes a 1% water emulsion.)

BUILDINGS HAVING BASEMENTS OR CRAWL SPACES

Dig a trench along the outside of the foundation walls. In brick or hollow block or concrete foundations, dig a trench to, but not below, the footing. Then as the trench is refilled, treat the soil at the rate of 4 gallons of above dilution per 10 linear feet for each foot of depth. A trench 3 feet deep would require 12 gallons per 10 linear feet.

The objective is to establish a treated soil barrier which will prevent termite entry. Other critical areas may also require treatment. It is suggested that you consult your State Extension Service for advice.

NOTICE: Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

MP-7K



CHLORDANE 8E

(8 lbs. Chlordane per Gallon)

AGRICULTURAL INSECTICIDE

ACTIVE INGREDIENTS

*Chlordane, technical 72.3%
Petroleum Distillate 21.5%

INERT INGREDIENTS 6.2%

*Equivalent to 43.38% octachloro-4,7-meth-
anotetrahydroindane and 28.92% related
compounds.

E.P.A. Reg. No. 802-71-AA

WARNING — Keep out of reach of children

See right side panel for additional warnings.

NET CONTENTS _____ GALLONS

THE CHAS. H. LILLY CO.
PORTLAND, OREGON 97218
EPA Est. 802-OR-1

LILLY'S

Miller's

ACCEPTED

DEC 10 1975

Under the Federal Insecticide, Fungicide, and Rodenticide Act

CHLORDANE 8E

(8 lbs. Chlordane per Gallon)
AGRICULTURAL INSECTICIDE

ACTIVE INGREDIENTS

*Chlordane, technical 72.3%
 Petroleum Distillate 21.5%

INERT INGREDIENTS

*Equivalent to 43.38% octachloro-4,7-meth-
 anotetrahydroindane and 28.92% related
 compounds.

E.P.A. Reg. No. 802-71-AA

WARNING — Keep out of reach of children
 See right side panel for additional warnings.

NET CONTENTS _____ GALLONS

THE CHAS. H. LILLY CO.
 PORTLAND, OREGON 97218
 EPA Est. 802-OR-1



Lilly

ACCEPTED
 DEC 10 1975
 Under the Federal Insecticide,
 Fungicide, and Rodenticide
 Act as amended, for the pesticide
 registered under the Pesticide
 Act, EPA Reg. No. 802-71

**PLEASE READ ENTIRE LABEL. USE ONLY AS DIRECTED,
 NOTING ALL WARNINGS, AND PRECAUTIONS**

WARNING

May be fatal if swallowed. Avoid breathing spray mist. Contact with skin can
 cause toxic symptoms. In case of contact with skin, wash with soap and water.
 Avoid contamination of feed and foodstuffs.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME

USE PRECAUTIONS

Do not apply or allow to drift to areas occupied by humans or animals.
 Do not apply to food crops after edible parts start to form. On Canberries,
 make application before blooming or after harvest. Consult State Agricultural
 Experiment Station or State Extension Service for exact timing and spacing of
 sprays in your area.

Do not feed treated forage to dairy animals or animals being finished for
 slaughter and do not allow these animals to graze on treated pasture, forage,
 range or wasteland. This product is highly toxic to bees exposed to direct
 treatment or residues on crops. Protective information may be obtained from
 your Cooperative Agricultural Extension Service. Do not apply when honey
 bees are foraging in the field.

Avoid drift to adjoining food and forage crops about to be harvested or crops
 in bloom. The grower is responsible for residue on his crops as well as for
 damages caused by drift from his property to that of others. Use only at
 recommended rates and observe use precautions in order that residues on
 food crops will not exceed established tolerances.

This product is toxic to fish and wildlife. Birds feeding on treated areas may
 be killed. Keep out of lakes, streams or ponds. Do not apply where runoff is
 likely to occur. Do not apply when weather conditions favor drift from areas
 treated. Do not contaminate water by cleaning of equipment or disposal of
 wastes. Apply this product only as specified on this label.

Rinse equipment and containers and dispose of wastes by burying in non-
 crop lands away from water supplies. Containers should be disposed of by
 breaking or punching holes in them and burying with wastes.

~~Use 1 to 2 pints per acre.~~
FLEA BEETLES—Beans, Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Kohlrabi

Use 1 to 2 pints per acre. Repeat applications if necessary.

ONION THRIPS—Onions

Use 1 to 2 pints per acre as a foliage spray. Repeat applications if necessary. Do not apply to green or spring onions.

ROOT MAGGOTS—Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Kohlrabi, Radishes, Rutabagas, Turnips

Use 1 to 2 pints per acre worked into top 3 or 4 inches of soil, as a pre-planting, planting time or transplanting treatment.

ANTS—Lawns, Ornamental Turfs, Parks, Cemeteries, Nursery Stock, Ornamentals.

Use 1 pint per acre. Apply directly on ants, along their trails and douse hills or nests.

Birds feeding on treated areas may be harmed.

Keep children and pets off treated areas until this material is washed into the soil and grass is dry.

TERMITES (Subterranean)

Use 4¼ qts. Miller's Chlordane 8E in 98-15/16 gals. of water. This makes a 1% water emulsion.)

BUILDINGS HAVING BASEMENTS OR CRAWL SPACES

Dig a trench along the outside of the foundation walls. In brick or hollow block or concrete foundations, dig a trench to, but not below, the footing. Then as the trench is refilled, treat the soil at the rate of 4 gallons of above dilution per 10 linear feet for each foot of depth. A trench 3 feet deep would require 12 gallons per 10 linear feet.

The objective is to establish a treated soil barrier which will prevent termite entry. Other critical areas may also require treatment. It is suggested that you consult your State Extension Service for advice.

NOTICE: Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

MP-7K