

LINDANE 25W

(25% Lindane Wettable Powder)

INSECTICIDE

**Livestock
Fly Control . . . Ornamentals**

Active ingredient	
Lindane (Gamma Isomer of Benzene Hexachloride)	25%
Inert Ingredients	75%
Total	100%

ACCEPTED
 Jul 24, 1978
 UNDER THE FEDERAL
 FERTILIZER ACT
 REGISTERED
 LINDANE 25W
 IS ATTACHED TO THIS



WARNING — Keep out of reach of children

May be fatal if swallowed! May be absorbed through skin! Do not breathe dust or spray mist. Avoid contact with skin and eyes. Wash thoroughly after handling. Avoid contamination of feed and foodstuffs.

E.P.A. Reg. No. 802-76-AA

Net wt. 4 lbs.

THE CHAS. H. LILLY CO.
 PORTLAND, OREGON

Miller's LINDANE 25W

DIRECTIONS FOR USE — LIVESTOCK SPRAYS

BEEF CATTLE—to control Lice or Ticks

1. Use at the rate of 1 lb. per 100 gal. of water.
2. Spray to assure thorough and complete coverage.

BEEF CATTLE—to control Mange Mites (Scab or Sarcoptic Mange)

1. Use at the rate of 1 ½ lbs. per 100 gal. of water.
2. Complete, thorough coverage of entire animal is essential to obtain the best results.
3. Make two applications only, one or two weeks apart.

SHEEP AND GOATS—to control Lice, Ticks and Scab Mites (Scabies)

1. As a Spray—Use 1 lb. per 100 gal. of water.
2. Apply to thoroughly wet animals.
3. For Scab control, make two applications 10 days apart.

SWINE—to control Lice and Sarcoptic Mange

1. Use 1 ½ lbs. per 100 gal. of water.
2. Spray thoroughly but do not over wet.

USE PRECAUTIONS — LIVESTOCK

1. Use only specified amount of chemical when mixing spray.
2. Make sure there is thorough and complete mixing by proper agitation before using spray solution.
3. Do not spray within 30 days of slaughter.
4. Do not spray animals under 3 months of age.
5. When necessary to spray young animals (over 3 months of age), use no more than 1 lb. per 100 gallons of water.
6. Do not treat sows within 2 weeks of farrowing nor for at least 3 weeks thereafter.
7. Do not use in poultry houses. Do not use on dairy cattle or on dairy goats.
8. CAUTION—as a general precaution in spraying animals, avoid treating during cold, stormy weather. Do not treat sick or emaciated animals.

FLY CONTROL (House and Stable Flies)

1. Residual Spray for Barns and Outbuildings—use 10 lbs. per 100 gal. of water. Apply as a coarse spray to thoroughly wet walls, windows, fences and other exterior surfaces to point of runoff.
2. Interior Spray for Barns, Outbuildings—use 10 lbs. per 100 gal. of water. Apply as a coarse spray to thoroughly wet walls, ceilings, beams, screens, and floors, using approximately 1 gallon diluted spray per 1,000 sq. ft.
3. Do not use in dairy barns or milk rooms. Do not contaminate animals' feed or water.

SYMPHYLID CONTROL ON ORNAMENTALS

1. Use at the rate of 4 oz. (of the dry powder) per 1000 sq. ft. and work thoroughly into the top 6 inches of soil. This will also aid in the control of root weevil larvae and white grubs.
2. Do not use on soil to be used for production of root or vegetable crops as lindane may cause an off-flavor.

USE PRECAUTIONS

This product is toxic to fish and wildlife. Keep out of lakes, ponds and streams. Do not apply to any area not specified on label.

Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes.

NOTICE: Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

