

Miller's  
**LINDANE 75W**

DOSAGE OF MILLER'S LINDANE 75W


for the temporary protection of  
germinating seeds and seedlings  
against Wireworms.

Seed	Ounces to be used per 100 lbs. of Seed
Berley . . . . .	1-1/3
Beans (Snap, Green, Wax) . . . . .	2/3 to 1-1/3
Beans (Lima) . . . . .	2/3
Beans (Large Lima) . . . . .	2/3 to 1-1/3
Beans (Fordhook Lima) . . . . .	1/3 to 2/3
Cantaloupes . . . . .	1-1/3
Corn . . . . .	1-1/3
Cucumbers . . . . .	1-1/3
Lettuce . . . . .	1-1/3
Milo . . . . .	1-1/3
Peas . . . . .	1-1/3
Sugar Beets . . . . .	4
Tomatoes . . . . .	2-2/3
Wheat . . . . .	1/3 to 2/3

**DRY TREATMENT** — Mix the required amount of Miller's LINDANE 75W with the seed in any commercial or barrel treater, until thoroughly coated.

**SLURRY TREATMENT** — Add the required amount of Miller's LINDANE 75W to the fungicide-slurry. One pint of liquid should be used for each 100 lbs. of Sugar Beet seed; 1/2 pint of liquid slurry to each 100 lbs. of Bean, Pea, Cucumber, Tomato, or Corn seed. Slurry treated seeds should not be stacked until the seed has had an opportunity to dry. If seed is stacked when wet, heat is developed which impairs the germination.

- NOTE . . .**
1. Treated seeds should be stored in a dry place until planted so that excess moisture may evaporate.
  2. Weigh accurately. Use only amounts suggested.
  3. Plant treated seed within 3 weeks time.
  4. Never permit treated seed to be used as food by humans or domestic animals.
  5. Bags containing treated seed should be so marked and should not be reused for any other purpose.



**LINDANE  
75W**

(75% Lindane, Wettable Powder)

**SEED TREATMENT  
WITH DYE  
for**

THE TEMPORARY PROTECTION OF  
GERMINATING SEEDS AND  
SEEDLINGS AGAINST

**WIREWORMS**

Active ingredient  
Gamma isomer of Benzene  
Hexachloride from Lindane . . . . . 75%  
Inert ingredients . . . . . 25%

**WARNING — Keep out of reach of children**

See right side panel for additional warnings.

E.P.A. Reg. No. 802-81-AA

Net Weight 1 Lb.

— MILLER PRODUCTS —  
**THE CHAS. H. LILLY CO.**  
PORTLAND, OREGON 97214

Miller's  
**LINDANE 75W**

Miller's LINDANE 75W may be used as temporary protection of germinating seed and seedlings of certain vegetable and field crops by means of seed treatment. It can be used with most of the common fungicides such as Hexachlorobenzene, Arasan and may be applied by the slurry method or in the dry form.

Use only amounts suggested — higher concentrations may cause seed injury. Plant treated seed within 3 weeks of treating, or delayed sprouting may result with some of the sensitive varieties.

Consult State Agricultural Experiment Station or State Extension Service Authorities for specific information for your area.

**WARNING**

May be Fatal if Swallowed! Do not breathe dust or spray mist. May be Absorbed Through Skin! Avoid contact with skin and eyes. Wash thoroughly after handling. Work in a well ventilated room.

Avoid contamination of feed and foodstuffs. Treated seed should be labeled as follows:

TREATED SEED. DO NOT USE FOR FOOD,  
FEED OR OIL PURPOSES.

This product is toxic to fish and wildlife. Keep out of lakes, streams, or ponds. Treated seeds on soil surface may be hazardous to birds and other wildlife. Dispose of excess treated seeds by burial in non-crop lands away from water supplies. Do not contaminate water by cleaning of equipment or disposal of wastes. Use this product only as specified on this label.

Rinse equipment and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes.

**NOTICE:** Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

802-81

8-25-72

1 of 2

