

GENERAL DIRECTIONS

1. Spray on first appearance of insect and repeat as necessary. For Red Spider Mites make at least two applications at 5 to 7-day intervals.
2. Spray to thoroughly cover plant or tree, including underside of leaves.
3. On Vegetables apply spray at the rate of 1 gal. to 300 sq. ft. garden area; or 3 gal. to 900 sq. ft.

ROSES, ORNAMENTALS, FLOWERS

1. For Green Aphids . . . use 1 teaspoon to 1 gallon of water; or 1 tablespoon to 3 gallons; or 1 pint to 100 gallons.
2. For Black Aphids, Red Spider Mites, Leaf Hoppers, Mealy Bugs, Thrips, Tarnished Plant Bugs, White Fly . . . use 2 teaspoons to 1 gallon of water; or 2 tablespoons to 3 gallons; or 1 pint to 50 gallons.
3. For soft Brown Scale, young crawler Scale . . . use 1 tablespoon to 1 gallon of water; or 3 tablespoons to 3 gallons; or 3 pints to 100 gallons.
4. Injury may occur on certain ferns, including Boston, Maidenhair and Pteris, as well as some species of Crassula.

EVERGREENS

1. For Red Spider Mites . . . use 2 teaspoons to 1 gallon of water or 2 tablespoons to 3 gallons; or 1 pint to 50 gallons.
2. For Monterey Pine Scale . . . use 1 tablespoon per gallon of water or 3 tablespoons to 3 gallons; or 3 pints to 100 gallons.

VEGETABLES

Beans, Cucumbers, Melons, Squash, Tomatoes (1*); Broccoli, Peas, Turnips (3*); Brussels Sprouts, Cabbage, Cauliflower, Kale, Mustard (7*); Potatoes (0*). (*The numbers indicate the number of days waiting time between spraying and harvest (for example, spray Tomatoes at 9:00 A.M. Monday; pick them after 9:00 A.M. Tuesday.)

1. For Green Aphids . . . use 1 teaspoon to 1 gallon of water; or 1 tablespoon to 3 gallons; or 1 pint to 100 gallons.
2. For Black Aphids, Red Spider Mites, Leaf Hoppers, Cabbage Loop, Imported Cabbage Worm, Mexican Bean Beetle . . . use 2 teaspoons to 1 gallon of water; or 2 tablespoons to 3 gallons; or 1 pint to 50 gallons.

FRUITS

Apples (3*); Pears (1*) (*The numbers indicate the number of days waiting time between spraying and harvest.)

1. For Aphids, Red Spider Mites, Pear Psylla . . . use 2 teaspoons to 1 gallon of water; or 2 tablespoons to 3 gallons; or 1 pint to 50 gallons.

NET CONTENTS
ONE QUART

\$5.98



Garden Malathion

57% Premium Grade Malathion

APHIDS, LEAF HOPPERS,
RED SPIDER MITES,
THRIPS, MEALY BUGS
FLOWERS, EVERGREENS,
VEGETABLES

ACCEPTED

AUG 27 1973

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND
FOR EC
ED T

CAUTION Keep out of the reach of children
See right side panel for additional cautions

E.P.A. Reg. No. 802-123-AA

Please read entire label.

Use only as directed, noting all cautions.

FLY CONTROL — OUTDOORS . . . Residual Spray

1. Use 4 teaspoons Miller's Garden MALATHION plus 2 tablespoons sugar to 1 quart of water; or 5 tablespoons Miller's Garden MALATHION plus 1/2 cup sugar to one gallon water; or 1 cup Miller's Garden MALATHION plus 1 cup sugar to 2 1/2 gallons of water.
2. Spray on walls, fences, garbage containers and wherever flies congregate and breed outdoors.
3. Wet all surfaces thoroughly. Repeat as necessary.
4. Do not allow this spray dilution to contact ornamental plants as injury may occur. Do not allow spray mist to contaminate fish ponds.

HOUSEHOLD INSECTS: Roaches, Ants, Spiders, Silverfish, Crickets, Clover Mites

1. Use 3 tablespoons to 1 quart of water; or 3/4 cup to 1 gallon.
2. Apply by means of a coarse spray, or paint brush, to window sills, baseboards, drainboards, under sinks, stoves, and to other areas frequented by insects. Repeat as necessary.
3. Water-based sprays may stain finished surfaces.
4. Do not contaminate food or utensils.

CAUTION . . . Harmful if inhaled, swallowed, or absorbed through the skin. Avoid breathing spray mist. Avoid contact with skin, eyes and clothing. Store away from children, domestic animals and foodstuffs. In case of contact, wash with soap and water; for eyes, get medical attention. Malathion upon use may cause cholinesterase inhibition. If poisoning occurs, get prompt medical aid. Note to Physician: Atropine is antidotal.

Cover or remove exposed foods. Avoid contamination of cooking utensils and food preparation surfaces.

This product is toxic to fish. Keep out of any body of water. Do not contaminate water by cleaning of equipment, or disposal of wastes. For the protection of honey bees, do not apply when they are foraging in the garden.

Destroy empty container. Do not re-use for any purpose.

NOTICE: Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith. MP:JHH

THE CHAS. H. LILLY CO.
PORTLAND, OREGON

802-123

8-27-73

1 of 3

2 of 3

PLEASE READ ENTIRE LABEL. USE ONLY AS DIRECTED, NOTING ALL CAUTIONS.

ROSES, ORNAMENTALS, FLOWERS

1. For Green Aphids . . . use 1 teaspoon per gallon of water.
2. For Black Aphids, Red Spider Mites, Leaf Hoppers, Mealy Bugs, Thrips, Tarnished Plant Bugs, White Fly . . . use 2 teaspoons per gallon of water.
3. For Soft Brown Scale, young crawler Scale . . . use 1 tablespoon per gallon of water.
4. Injury may occur on certain ferns including Boston, Maidenhair and Pteris, as well as some species of Crassula.

VEGETABLES

- Beans, Cucumbers, Melons, Squash, Tomatoes (1*); Broccoli, Peas Turnips (3*); Brussels Sprouts, Cabbage, Cauliflower, Kale, Mustard (7*); Potatoes (0*). (*The numbers indicate the number of days waiting time between spraying and harvest (for example, spray Tomatoes at 9:00 A.M. Monday; pick them after 9:00 A.M. on Tuesday.))
1. For Green Aphids . . . use 1 teaspoon per gallon of water.
 2. For Black Aphids, Red Spider Mites, Leaf Hoppers, Cabbage Looper, Imported Cabbage Worm, Mexican Bean Beetle . . . use 2 teaspoons per gallon of water.

NET CONTENTS
8 FL. OZ.

\$2.29



Garden

Malathion

57% Premium Grade Malathion



UNDER THE INSECT FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTERED UNDER NO. 723 SUBJECT TO ATTACHED COMMENTS.

CAUTION Keep out of the reach of children
See right side panel for additional cautions
E.P.A. Reg. No. 802-123-AA

GENERAL DIRECTIONS

1. Spray on first appearance of insect and repeat as necessary. For Red Spider Mites make at least two applications at 5 to 7-day intervals.
2. Spray to thoroughly cover plant or tree, including underside of leaves.
3. On Vegetables, apply spray at the rate of 1 gal. to 300 sq. ft. of garden area.

CAUTION

Harmful if inhaled, swallowed or absorbed through the skin. Avoid breathing spray mist. Avoid contact with skin, eyes and clothing. Store away from children, domestic animals and foodstuffs. In case of contact, wash with soap and water; for eyes, get medical attention. Malathion may cause cholinesterase inhibition. If accidental poisoning occurs, get prompt medical aid. **Note to Physician:** Atropine is antidotal.

Cover or remove exposed foods. Avoid contamination of cooking utensils and food preparation surfaces. This product is toxic to fish. Keep out of any body of water. Do not contaminate water by cleaning of equipment, or disposal of wastes. For the protection of honey bees, do not apply when they are foraging in the garden. Destroy empty container. Do not re-use for any purpose.

Active ingredients	
*Malathion	57%
Aromatic Petroleum	
Derivative Solvents	33%
Inert ingredients	10%
*O-O-dimethyl dithiophosphate of diethyl mercaptosuccinate.	

NOTICE: Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith. MP-11H

THE CHAS. H. LILLY CO.
PORTLAND, OREGON

GENERAL DIRECTIONS

1. Spray on first appearance of insect and repeat as necessary. For Red Spider Mites make at least two applications at 5 to 7-day intervals.
2. Spray to thoroughly cover plant or tree, including underside of leaves.
3. On Vegetables, apply spray at the rate of 1 gal. to 300 sq. ft. of garden area; or 3 gal. to 900 sq. ft.

ROSES, ORNAMENTALS, FLOWERS

1. For Green Aphids . . . use 1 teaspoon per gallon of water.
2. For Black Aphids, Red Spider Mites, Leaf Hoppers, Mealy Bugs, Thrips, Tarnished Plant Bugs, White Fly . . . use 2 teaspoons per gallon of water.
3. For Soft Brown Scale, young crawler Scale . . . use 1 tablespoon per gallon of water.
4. Injury may occur on certain ferns, including Boston Maidenhair and Pteris, as well as some species of Crassula.

EVERGREENS

1. For Red Spider Mites . . . use 2 teaspoons per gallon of water.
2. For Monterey Pine Scale . . . use 1 tablespoon per gallon of water.

VEGETABLES . . .

Beans, Cucumbers, Melons, Squash, Tomatoes (1*); Broccoli, Peas, Turnips (3*); Brussels Sprouts, Cabbage, Cauliflower, Kale, Mustard (7*); Potatoes (0*). (*The numbers indicate the number of days waiting time between spraying and harvest (for example, spray Tomatoes at 9:00 A.M. Monday; pick them after 9:00 A.M. on Tuesday.)

1. For Green Aphids . . . use 1 teaspoon per gallon of water.
2. For Black Aphids, Red Spider Mites, Leaf Hoppers, Cabbage Looper, Imported Cabbage Worm, Mexican Bean Beetle . . . use 2 teaspoons per gallon of water.

NET CONTENTS
ONE PINT

\$3.69



Garden Malathion

57% Premium Grade Malathion



CAUTION Keep out of the reach of children
See right side panel for additional cautions
E.P.A. Reg. No. 802-123-AA

PLEASE READ ENTIRE LABEL. USE ONLY AS DIRECTED, NOTING ALL CAUTIONS.

FLY CONTROL—OUTDOORS . . . Residual Spray

1. Use 4 teaspoons Miller's Garden MALATHION plus 2 tablespoons sugar to 1 quart of water.
2. Spray on walls, fences, garbage containers and wherever flies congregate and breed outdoors.
3. Wet all surfaces thoroughly. Repeat as necessary.
4. Do not allow this spray dilution to contact ornamental plants as injury may occur. Do not allow spray mist to contaminate fish ponds.

Active ingredients

*Malathion 57%
Aromatic Petroleum Derivative Solvents 33%

Inert ingredients 10%

*O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate

CAUTION

Harmful if inhaled, swallowed, or absorbed through the skin. Avoid breathing spray mist. Avoid contact with skin, eyes and clothing. Store away from children, domestic animals and foodstuffs. In case of contact, wash with soap and water; for eyes, get medical attention. Malathion upon use may cause cholinesterase inhibition. If poisoning occurs, get prompt medical aid. Note to Physician: Atropine is antidotal.

Cover or remove exposed foods. Avoid contamination of cooking utensils and food preparation surfaces.

This product is toxic to fish. Keep out of any body of water. Do not contaminate water by cleaning of equipment, or disposal of wastes. For the protection of honey bees, do not apply when they are foraging in the garden.

Destroy empty container. Do not re-use for any purpose.

NOTICE: Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith. MP-11H

THE CHAS. H. LILLY CO.
PORTLAND, OREGON

APC
W