

Miller's 2,4,5-T AMINE 4

(4 lbs. 2,4,5-T acid equivalent per gallon)

GENERAL DIRECTIONS

Add half the required amount of water to the spray tank, then add the specified amount of Miller's 2,4,5-T AMINE 4 with agitation, then add the balance of the water with continued agitation. Mix thoroughly.

Do not graze dairy animals on treated areas within 6 weeks after application. Do not graze meat animals on treated areas within 2 weeks of slaughter.

SPECIFIC DIRECTIONS

FOLIAGE TREATMENT

1. Use 3 quarts per 100 gallons of water.
2. For woody growth up to 6 to 8 feet tall, apply after foliage is well developed and plants are actively growing, using a drenching spray.
3. Brush over 6 or 8 feet tall can be sprayed successfully, although in many cases Basal Bark or Stump Treatment is preferable.
4. Thorough coverage is important, wetting all parts of the plants, including foliage, shoot stems and bark.
5. Best results usually will be obtained from applications made soon after maximum foliage development in the spring. With good growing conditions, applications may be made up to 2 to 3 weeks before frost time. Power equipment with pressures up to 250 pounds will aid in obtaining satisfactory spray coverage. Repeat applications may be required as new growth develops. Usually one good treatment a year should be adequate.

BASAL BARK TREATMENT

1. Use 4 gallons in 96 gallons of water (1 pint in 3 gallons).
2. Spray the basal parts of brush stems and tree trunks to a height of 12 to 15 inches from the ground line, using low pressures.
3. Apply the spray until run-off to the ground line is noticeable.
4. More spray volume is necessary for old or rough bark than for young or smooth bark.
5. Applications may be made at any time including the winter months.
6. Treated plants should not be cut for one full season after treatment.
7. Some plants can be controlled with a solution of 2 to 3 gallons in 97 to 98 gallons of water.

Miller's 2,4,5-T AMINE 4

SPECIFIC DIRECTIONS

STUMP TREATMENT

1. Use 4 gallons in 96 gallons of water (1 pint in 3 gallons).
2. Cut growth more than 6 to 8 feet tall close to the ground, then spray the stumps and stubs.
3. Wet thoroughly all exposed bark and cut surfaces as described under Basal Bark treatment.

GIRDLE TREATMENT

1. Use 2 gallons in 100 gallons of water (½ pint in 3 gallons).
2. Make a single girdle of overlapping axe cuts, at a convenient height, on large trees.
3. Apply the spray thoroughly to the injured area.

Consult State Agricultural Experiment Station or State Extension Service Weed Specialists for specific advice as climate and local conditions may cause suggested uses to vary.



2,4,5-T AMINE 4

(4 lbs. 2,4,5-T acid equivalent per gallon)

For the control of Herbaceous and Woody Plants growing in Pasture, Rangeland, Fence Rows, Farmyards and Right-of-Ways

| | | |
|--------------|-------------|-----------------|
| Ash | Maple | Wild Blackberry |
| Brambles | Oak | Wild Cherry |
| Chokecherry | Poison Oak | Wild Rose, |
| Hawthorn | Poison Ivy | Certain Species |
| Horse Nettle | Salmonberry | of Ribes |

Active ingredient
Triethylammonium
2,4,5-Trichlorophenoxyacetate 57.2%
Inert ingredients 42.8%
Total 100.0%
*Equivalent to 41.1% 2,4,5-Trichlorophenoxyacetic acid.

E.P.A. Reg. No. 802-128-AA

BRUSH KILLER

CAUTION — Keep out of reach of children

See right side panel for additional cautions

1 GALLON

ACCEPTED
UNDER THE FEDERAL
FUNGICIDE AND ROX
FOR PESTICIDE POS
EPA REG. NO. 112

— MILLER PRODUCTS —
THE CHAS. H. LILLY CO.
PORTLAND, OREGON 97214

Miller's
**2,4,5-T
AMINE 4**

SPECIFIC DIRECTIONS

STUMP TREATMENT

1. Use 4 gallons in 96 gallons of water (1 pint in 3 gallons).
2. Cut growth more than 6 to 8 feet tall close to the ground, then spray the stumps and stubs.
3. Wet thoroughly all exposed bark and cut surfaces as described under Basal Bark treatment.

GIRDLE TREATMENT

1. Use 2 gallons in 100 gallons of water (½ pint in 3 gallons).
2. Make a single girdle of overlapping axe cuts, at a convenient height, on large trees.
3. Apply the spray thoroughly to the injured area.

Consult State Agricultural Experiment Station or State Extension Service Weed Specialists for specific advice as climate and local conditions may cause suggested uses to vary.



**2,4,5-T
AMINE 4**

(4 lbs. 2,4,5-T acid equivalent per gallon)

For the control of Herbaceous and Woody Plants growing in Pastures, Rangeland, Fence Rows, Farmyards and Right-of-Ways

| | | |
|--------------|-------------|-----------------|
| Ash | Maple | Wild Blackberry |
| Brambles | Oak | Wild Cherry |
| Chokecherry | Poison Oak | Wild Rose, and |
| Hawthorn | Poison Ivy | Certain Species |
| Horse Nettle | Salmonberry | of Ribes |

| | |
|---|--------|
| Active ingredient | |
| Triethylammonium | |
| 2,4,5-Trichlorophenoxyacetate | 57.2% |
| Inert ingredients | 42.8% |
| Total | 100.0% |
| *Equivalent to 41.1% 2,4,5-Trichlorophenoxyacetic acid. | |


E.P.A. Reg. No. 802-128-AA

BRUSH KILLER

CAUTION — Keep out of reach of children

See right side panel for additional cautions

1 GALLON



— MILLER PRODUCTS —
THE CHAS. H. LILLY CO.
PORTLAND, OREGON 97214



CAUTION

PLEASE READ ENTIRE LABEL. USE ONLY AS DIRECTED, NOTING ALL CAUTIONS AND WARNINGS.

Harmful if swallowed.

This product may cause skin irritation. Avoid contact with skin, eyes and clothing. In case of contact, wash skin with plenty of soap and water; flush eyes with plenty of water for at least 15 minutes and get medical attention. Keep out of reach of children and domestic animals.

Do not use around the home or recreation areas.

Do not store with foodstuffs. Do not transport or store near fertilizers, insecticides, fungicides or seeds.

Do not apply directly to or otherwise permit sprays to come in contact with vegetables, flowers, grapes, fruit trees, cane berries, ornamentals or other desirable plants which are sensitive to 2,4-D and do not permit spray mist to drift onto them, since even minute quantities of the spray may cause severe injury during both growing and dormant periods. Coarse sprays are less likely to drift.

Clovers may be injured with this material.

Application by airplane, ground equipment, or hand sprayers should be made only when there is no hazard from drift.

Do not apply by airplane in the vicinity of grapes or other desirable 2,4,5-T susceptible plants.

Do not contaminate irrigation ditches or water used for domestic purposes. Excessive amounts of 2,4,5-T in the soil may temporarily inhibit seed germination or plant growth.

At higher temperatures vaporization may cause injury to susceptible plants growing nearby. Keep container tightly closed when not in use.

Do not use spray equipment for any other purpose. Such equipment should not be used for application of insecticides, fungicides, or other agricultural chemicals.

Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes. Do not contaminate water by cleaning of equipment, or disposal of wastes.

NOTICE: Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

MP 5G