

**PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS AND DOMESTIC ANIMALS**

CAUTION

Harmful if swallowed. Avoid contact with skin, eyes or clothing. Avoid breathing spray mist. Wash thoroughly after handling.

STATEMENT OF PRACTICAL TREATMENT

In case of contact, wash skin with soap and water; for eyes flush with water for 15 minutes and get medical attention.

ENVIRONMENTAL HAZARD

Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply when weather conditions favor drift from target area.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

LILLY/MILLER
Dandelion Killer



ACTIVE INGREDIENT	
*2,4-Dichlorophenoxyacetic Acid Dimethylamine Salt	13.13
INERT INGREDIENTS	86.87
Total	100.00
*2,4-Dichlorophenoxyacetic acid equivalent 10.9%†	
†Isomer Specific by AOAC method No. 6.001-5	

Kills Dandelions, Chickweed, Knotweed, Plantains and other broadleaf weeds in lawns.

Keep out of reach of children
CAUTION

See side panels for Statement of Practical Treatment and additional Precautionary Statements.

NET CONTENTS 1 QUART

STORAGE AND DISPOSAL

STORAGE: Protect from freezing. Do not store near feed or food stuffs.
DISPOSAL: Do not reuse empty container. Wrap container and put in trash collection.

NOTICE: Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given hereon.

LC-1R

THE CHAS. H. LILLY CO.
PORTLAND, OREGON 97218

EPA Est. 802-OR-1
EPA Reg. No. 802-148

SHAKE WELL BEFORE USING.

LAWN WEED CONTROL

Spring (April-May) and Fall (September-October) applications for:

Chickweed	Henbit	Plantains
Dandelions	Knotweed	Thistles
Docks	Morningglory	...and other broadleaf weeds.

Amount to use... Established Lawns

Use 2 tablespoons per gallon of water per 200 sq. ft. of lawn. Wet solid patches of weeds thoroughly.

New Lawns

Use 1 tablespoon per gallon of water per 200 sq. ft. of lawn. Do not use until lawn is well established ... after second or third cutting

NOTE: Do not use on St. Augustine grass, Dichondra or Carpetgrass lawns. Bentgrass and Clover lawns may be injured with the material.

SPRAYING DIRECTIONS

1. Cover area uniformly with a coarse droplet spray, when there is little or no wind, so that the spray solution will not drift to susceptible vegetation such as flowers, vegetables, grapes, fruit trees and ornamentals.

2. Apply with a hose-end sprayer or a pressure sprayer designed for lawns. (Use a separate sprayer for weed killing only)

HOSE-END SPRAYER: Put recommended amount of product in bottle, add water to proper level and spray area indicated following chart below. For example, to spray 1600 sq. ft. of lawn area, measure 16 tablespoons into jar, add water to the 8 gallon level and mix thoroughly. Attach jar to hose, turn water on and spray as evenly as possible over the 1600 sq. ft.

Amount Lilly/Miller Dandelion Killer to use:

TBS.	FL. OZ.	Fill to Gallon(s) Level	Area to Treat (Sq. Ft.)
2	1	1	200
8	4	4	800
16	8	8	1600
32 (1 pint)	16	16	3200

PRESSURE SPRAYERS: Measure recommended amount of product into tank and dilute with water according to the following chart:

TBS.	FL. OZ.	Amount of Water (Gals)	Area to Treat
2	1	1	200
4	2	2	400
10	5	5	1000
32 (1 pint)	16	16	3200
64 (1 quart)	32	32	6400

SPOT APPLICATION

For Chickweed, False Dandelion and certain other hard-to-kill weeds.

1. Use 4 tablespoons per gallon of water in pressure sprayer (do not use hose end sprayer)
2. "Spot" spray small area or individual weeds, being certain to wet each weed.

APPLICATION TIPS

For best results, weeds should be actively growing with enough leaf area to absorb the chemical.

1. Do not cut lawn 2 to 3 days before treating
2. Soil should be moist at the time of application
3. Wet solid patches of weeds thoroughly

For best results, allow the chemical to move throughout the weed.

1. Withhold water 1 to 2 days after treating
2. Do not mow the lawn for 2 to 3 days after treating