

## DIRECTIONS

### PREVENTION IS THE KEYNOTE TO FUNGUS DISEASE CONTROL

#### VEGETABLES

**GENERAL** . . . Damping-off . . . see Seed Bed Treatment

**TOMATOES, POTATOES** . . . Late Blight

1. Dust or spray on first sign of disease, preferably before disease appears.
2. Repeat at 4 to 7-day intervals until under control.

**POTATOES—Seed Piece Treatment** . . . Seed Decay (Fusarium coeruleum)

1. Use 1 tablespoon per 2 pounds cut seed pieces.
2. Place dust and pieces in bag and shake thoroughly until seed pieces are completely covered.
3. Plant immediately before pieces dry out.

#### FRUIT

**APPLES** . . . Scab

1. Use as a spray (see dilution above).
2. Apply in pre-blossom, petal fall and first cover sprays.
3. Repeat if infection persists.

**PEACHES, CHERRIES** . . . Brown Rot

1. Use as a spray (see dilution above).
2. Apply in pink or popcorn stage, full bloom, and petal or shuck fall stage.

**LAWNS, TURF** . . . Brown Patch, Damping-Off

1. **Dust** . . . 1 pound evenly over 350 sq. ft. lawn area and water thoroughly after treatment . . . or
2. **Spray** . . . use 1 lb. to 5 gal. of water (keep agitated) and apply 1 gallon to each 100 sq. ft. lawn area.
3. Apply before disease appears. Repeat at 7 to 10-day intervals during summer.

## CAUTION

Avoid inhalation of dust. Avoid contact with skin.

Avoid contamination of feed and foodstuffs.

This product is toxic to fish. Keep out of any body of water.

Do not contaminate water by cleaning of equipment, or disposal of wastes.

Destroy empty container. Do not re-use for any purpose.

NET WT.  
1 LB.

\$1.79



Garden

# Captan

(Garden Fungicide) Dust or Spray

#### Active ingredient

Captan (N[(trichloromethyl) thio]-4-cyclohexene-1,2-dicarboximide) 5%

Inert ingredients . . . . . 95%

Total . . . . . 100%

**CAUTION** Keep out of the reach of children  
See left side panel for additional cautions.

E.P.A. Reg. No. 802-152-AA

## DIRECTIONS

#### DUSTING

1. Dust lightly before diseases appear.
2. Repeat light dustings at 4 to 7-day intervals until under control.
3. Frequent light dustings (about 1 lb. per 800 sq. ft.) are more effective than single heavy dustings.

#### SPRAYING

1. Use 3 level tablespoons per quart of water; or 1 lb. per 5 gal.
2. Make thin paste with small amount of water, and then add balance of water required.
3. Spray lightly at 4 to 7-day intervals until under control.
4. Be sure underside of leaves is covered.

#### FLOWERS . . . ORNAMENTALS

**ROSES** . . . Black Spot

**CARNATIONS** . . . Rust, Leaf Spot (Alternaria)

**CHRYSANTHEMUMS** . . . Flower Blight (Botrytis)

1. Dust or spray on first sign of disease, preferably before disease appears.
2. Repeat at 4 to 7-day intervals until under control.

#### DAMPING-OFF

**AZALEA, CARNATION, and CHRYSANTHEMUM CUTTINGS**

1. Use 3 level tablespoons per pint of water.
2. Dip cuttings in this suspension before planting.
3. Stir diluted suspension before dipping to insure uniformity.

**SEED BED** . . . Soil Treatment, Before Planting

1. Dust 1 pound evenly over 350 sq. ft. of soil area.
2. Mix thoroughly with top 3 to 4 inches of soil.

#### AFTER SEEDLINGS EMERGE

1. Dust emerging seedlings lightly at 3 to 4-day intervals until plants are well established.

#### BULBS

**GLADIOLI, TUBEROUS BEGONIAS** . . . Rot (helpful in reducing certain rots)

1. Dust clean bulbs thoroughly before planting.

Captan . . . U.S. Patents No. 2,553,770; 2,553,771, 2,553,776.

**NOTICE:** Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith. MP-5F

**MILLER PRODUCTS**  
Portland, Oregon 97218