

LILLY/MILLER CAPTAN
GARDEN FUNGICIDE DUST OR SPRAY

ACTIVE INGREDIENT:

- *Captan5.0%
- Related derivatives0.1%
- INERT INGREDIENTS94.9%
- *N-Trichloromethylthio-4-cyclohexene-1,2-dicarboximide

FOR FLOWERS, ORNAMENTALS, BULBS, VEGETABLES, FRUITS, LAWNS.

Keep out of reach of children
DANGER
See side panel for additional caution

ACCEPTED
Per self-cert
OCT 26 1987

NET WEIGHT 1 LB.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER: Causes irreversible eye damage. Harmful if swallowed or inhaled. May cause allergic skin reactions. Do not get in eyes. Wear goggles or face shield when handling. Avoid contact with skin and clothing. Remove and separately launder clothing before reuse.

If In Eyes: Flush with plenty of water. Call a physician. If Swallowed: Drink promptly a large quantity of milk, egg whites, gelatin solutions, or if these are not available, drink large quantities of water. Avoid alcohol. If Inhaled: Remove victim to fresh air. If not breathing give artificial respiration, preferably mouth-to-mouth. Get medical attention.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

USE PRECAUTIONS

Wear goggles or face shield when handling. Remove contaminated clothing and separately launder clothing before reuse.

Wear chemical resistant gloves when using indoors or outdoors. When using wear long pants and long-sleeved shirt and apply with the wind to your back. Wash non-disposable gloves thoroughly with soap and water before removing.

HOW TO USE CAPTAN - Application Directions

Apply CAPTAN either by sprinkling the dust from container or mixing with water in a sprayer and spraying. To dust: 1) Make a light application and repeat every 4 to 7 days as necessary. 2) Get best control by making several light applications rather than one heavy application. To spray: 1) Use 3 level tablespoons per quart of water or 1 lb. per 5 gals. 2) Add a small amount of water, mix into a paste, then add the rest of the water. 3) Spray lightly, covering both leaf surfaces, every

BEST AVAILABLE COPY

4 to 7 days as necessary.

ORNAMENTALS

Foliage treatment for ROSES: Black Spot, CARNATIONS: Rust, Leaf Spot (Alternaria); CHRYSANTHEMUMS: Flower Blight (Botrytis). Dust or spray at first growth or sign of disease. See application directions above. Repeat every 4 to 7 days, until disease is under control.

Dip for cuttings of AZALEA, CARNATION, CHRYSANTHEMUM to control Damping Off: Make dip solution by using 3 level tablespoons per quart of water. Stir solution and dip cuttings into solution immediately before planting.

Preplant Dust Treatment for GLADIOLI, TUBEROUS BEGONIA to reduce rots: Add CAPTAN to paper bag. Place bulbs in paper bag, fold to close and gently shake bag until bulbs are coated with dust.

PREPLANT SOIL TREATMENT FOR VEGETABLE AND FLOWER SEED BEDS - Damping Off: Sprinkle 1 lb. CAPTAN per 350 sq. ft. over soil area prepared for planting. Mix dust thoroughly into the top 3 to 4 inches of soil. AFTER SEEDLINGS EMERGE, dust lightly every 3 to 4 days until plants are well established.

VEGETABLES

Foliage treatment for TOMATOES, to control Late Blight: Make a preventative treatment, then dust or spray with CAPTAN when disease appears. See Application Directions. Repeat applications every 4 to 7 days until disease is under control.

Seed Piece Treatment for POTATOES to control Seed Decay (Fusarium coeruleum): Use 1 tablespoon per 2 lbs. of cut seed pieces. Put CAPTAN and seed pieces in a bag, fold to close and shake bag until pieces are evenly coated with dust. Plant pieces immediately after treatment.

FRUIT

Spray applications for APPLES to control Scab: Mix spray solution according to Application Directions given above. Apply in pre-blossom, petal fall and first cover sprays. Repeat if infection persists.

PEACHES, CHERRIES: Brown Rot; See Application Directions above. Apply in pink or popcorn stage, full bloom, and petal or shuck fall stage.

LAWNS

Brown Patch, Damping Off: Dust or spray when growth appears in the ~~spring and repeat every 7 to 10 days during the spring and summer.~~

DUST: Apply 1 lb. CAPTAN per 350 sq. ft. Water thoroughly after application. SPRAY: Add 1 lb. to 5 gals. of water and apply 1 gal. per 100 sq. ft. of lawn area. Continuously agitate solution while spraying. Apply before disease appears and repeat every 7 to 10 days during the summer.

STORAGE AND DISPOSAL

Store this product in its original container and keep in a locked storage area out of reach of children and domestic animals. Do not reuse container. Securely wrap container in several layers of newspaper and discard in trash.

Filled by weight not by volume. Product may settle after packaging.

NOTICE: Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of

2063

this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

THE CHAS. H. LILLY CO.
PORTLAND, OREGON 97218
EPA Est. 802-OR-1
EPA Reg. No. 802-152
28603

Revised 10-6-87