



Miller's METHOXYCHLOR 5D

ACCEPTED
802-167
SEP 18 1973

(5% Methoxychlor Dust)

Active Ingredients

*Methoxychlor, technical
Inert Ingredients
*Equivalent to 4.4% 2,2-bis(4-chloroethoxy)propane and 0.6% other isomers and reaction products.

UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTERED UNDER NO. 802-167 SUBJECT TO ATTACHED COMMENTS (1)

CAUTION - Keep out of reach of children.

Avoid dust drift to areas occupied by unprotected humans or beneficial animals.

Wash treated produce thoroughly before eating or processing.

This product is toxic to fish. Keep out of lakes, ponds, or streams. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

Do not feed the treated crop or crop residue to poultry, dairy or meat animals.

Destroy empty containers. Bury wastes and empty containers in non-crop lands away from water, supplies or farm empty containers (stay away from snakes and fumes).

DIRECTIONS FOR USE

Apply on first appearance of insect or insect injury. Repeat applications as needed for adequate control.

Consult State Agricultural Experiment or State Extension Service authorities for information regarding timing of applications in your area.

Apply when there is little or no wind.

Crop and Pest	#Days	Pounds per acre
ALFALFA - Alfalfa Weevil Larvae, Alfalfa Caterpillar		
	7	30 to 40
CHEERRIES (For fresh market)		
Cherry Fruit fly	7	30
STRAWBERRIES - Spittle Bugs, Omnivorous Leaf Tier (Strawberry Fruit Worm)		
	14	45
ASPARAGUS - Asparagus Beetle, Flea Beetle		
	3*	45
*3 days unless washed or blanched.		
BEANS (Snap and Lima)		
Western Spotted Cucumber Beetle	3	40
CABBAGE, BROCCOLI, BRUSSELS SPROUTS		
Imported Cabbage Worm, Flea Beetles	14*	45
*3 days for Cabbage.		
CUCUMBERS, CANTALOUPES, WATERMELONS		
Striped Cucumber Beetle	1	35
Apply when plants are dry.		
LETTUCE - Western Spotted Cucumber Beetle		
	14	25
SPINACH - Western Spotted Cucumber Beetle		
	14	45
TOMATOES - Tuber Flea Beetle, Western Spotted Cucumber Beetle		
	1	25
TURNIPS, RUTABAGAS - Cabbage Flea Beetle		
	7*	45
*14 days if tops are to be used as food or feed.		

*Days indicate the number of days between last application and harvest.

E.P.A. Reg. No. 802-167-AA

17-31

THE CHAS. H. MILLER CO.
PORTLAND, OREGON

