

Miller's

## PET, POULTRY and LIVESTOCK DUST

### DIRECTIONS

#### LIVESTOCK — Beef Cattle, Non-Lactating Dairy Animals, Hogs Lice

1. Make thorough application to animals.
2. Repeat application in 10 days and thereafter as needed.
3. For **Hogs**, the pens should also be thoroughly dusted.
4. Do not apply on non-lactating dairy animals within two weeks of freshening. Do not apply to calves under one month of age.

#### LIVESTOCK — Beef and Dairy Cattle Hornflies

1. Apply 4 tablespoons per animal
2. Apply on the back and neck of animals and repeat applications at 10 to 14-day intervals.
3. Do not apply to dairy cattle later than five hours before milking or during milking

### USE PRECAUTIONS

Avoid contamination of feed, feeding troughs, and watering troughs or fountains.

### CAUTION

Avoid breathing dust. Avoid contact with skin. Wash thoroughly after using. Avoid contamination of feed and foodstuffs. Keep out of reach of children.

This product is toxic to fish. Keep out of lakes, streams and ponds.

Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes.

**NOTICE:** Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

MP 67

Insist on...

Miller's

\$1.49



PET

POULTRY

and

LIVESTOCK

(including Dairy)

DUST

(4% Malathion Dust)

CAUTION

Keep out of reach of children

See left side panel for additional cautions.

Active Ingredient

Malathion (O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate) 4.0%

Inert Ingredients 96.0%

E P A Reg No 802170-AF

NET WEIGHT 1 LB.

MILLER PRODUCTS

PORTLAND, OREGON 97218



Miller's

## PET, POULTRY and LIVESTOCK DUST

### DIRECTIONS

#### PETS — Dogs and Cats

##### Fleas on Animals

1. Use sufficient dust to treat animals thoroughly.
2. To avoid unnecessary application, animal should be closely examined before retreating. If necessary, repeat treatments should not be more often than once weekly.
3. If animal has free run, treatment may have to be made every 2nd or 3rd week.
4. If animal is confined to an area free of fleas, one treatment may be sufficient.

##### Fleas in Animal Quarters

1. Apply at rate of 1 or 2 lbs. per 1000 sq. ft., using hand duster.
2. Treat kennels, pens, yards, lawns, under porches or house. Remove manure and debris before dusting.

#### POULTRY — Chickens, Ducks, Geese, Turkeys

For direct application in the control of Northern Fowl Mite, Poultry Lice and as a supplement to premise treatments for Chicken Red Mite

1. Dust individual birds by means of a shaker can, or rotary hand duster.
2. Repeat in 4 to 5 weeks or when necessary.

For application in poultry houses in the control of Northern Fowl Mite, Chicken Red Mite, Poultry Lice

1. Use 1 lb. per 50 to 60 sq. ft. Repeat as necessary.
2. Apply dust to litter, floor space, and to nooks, crannies, and adjacent areas.
3. Apply with rotary hand duster, puff duster, or by blowing from a can or other container.