

# Miller's SPRA-OIL

## USE DIRECTIONS

### SPRING AND FALL SPRAY

Young Crawler Scale, Soft Brown Scale on Holly, Camellias, Azaleas, Rhododendrons, Roses, broadleaved Evergreen Trees and Shrubs

1. Use 5 teaspoons Miller's SPRA-OIL per quart of water or 3 1/2 tablespoons per gallon or 1 quart per 10 gallons
2. Add Miller's SPRA-OIL to sprayer slowly while filled with water. Mix thoroughly
3. **First spray**—Spring—before growth starts
4. **Second spray**—Fall (about September)—after growth

### DORMANT SPRAY—(Combination Spray—see table below)

Aphids' eggs, Red Spider Mites, Scale (Soft Brown, Rose, Pear, Oyster Shell), Blister Mite, Pear and Apple Scale, Mealy Bugs, Roses, Deciduous Ornamentals, Apples, Pears, Prunes, cherries,

1. Use Combination Spray (see table below) mixed and mixed thoroughly
2. Apply in Early Spring just before buds break

### DORMANT COMBINATION SPRAY

Miller's SPRA OIL	Miller's POLYSUL	Water
1 quart	1 quart	10 gallons
1 pint	1 pint	5 gallons
1 cup	1 cup	2 1/2 gallons

NET CONTENTS  
ONE QUART



Miller's

# Spra-oil

FOR CONTROL  
OF  
**RED SPIDER MITES, MEALY BUGS  
CRAWLER SCALE, HOLLY SCALE  
SOFT BROWN SCALE**

ACTIVE INGREDIENT Petreum Oil 98%  
NET INGREDIENTS 99%  
Miller's is registered for the 92%  
Grade Unclassified

**CAUTION** Keep out of the reach of children.  
See right side panel for additional cautions.

E.P.A. Reg. No. 802-179-AA

# Miller's SPRA-OIL

PLEASE READ ENTIRE LABEL. USE ONLY AS DIRECTED,  
NOTING ALL CAUTIONS AND WARNINGS.

## CAUTION

Keep out of reach of children. If swallowed **do not** induce vomiting. Call physician immediately.

Never apply a combination oil and sulfur spray to a plant. Do not apply an oil spray to foliage until 30 to 60 days have elapsed after a sulfur application as injury to foliage or flowers will result. Do not spray when weather is excessively hot. Be sure plants are partially well watered before spraying.

Keep out of lakes, streams, or ponds. Do not use in or near water. Do not use in or near water. Do not use in or near water. Do not use in or near water.

Do not use in or near water.

**NOTICE**—This product is not intended for use on food crops. Do not use on food crops. Do not use on food crops. Do not use on food crops.

THE MILLER COMPANY  
P.O. Box 100  
Piquette, Michigan 48864

ACCEPTED  
802-179

# Miller's SPRA-OIL

## USE DIRECTIONS

### SPRING AND FALL SPRAY

Young Crawler Scale, Soft Brown Scale on Holly, Camellias, Azaleas, Rhododendrons, Roses, broadleaved Evergreen Trees and Shrubs.

1. Use 5 teaspoons Miller's SPRA-OIL per quart of water; or 6 1/2 tablespoons per gallon; or 1 quart per 10 gallons.
2. Add Miller's SPRA-OIL to sprayer slowly while filling with water. Mix thoroughly.
3. **First spray**—Spring—before growth starts.
4. **Second spray**—Fall (about September)—after growth.

### DORMANT SPRAY—(Combination Spray, see table below)

Aphids' eggs, Red Spider Mites, Scale (Soft Brown, Rose, San Jose, Oyster Shell), Blister Mite, Pear and Apple Scab, Moss on Roses, Deciduous Ornamentals, Apples, Pears, Prunes, Cherries.

1. Use Combination Spray (see table below). Cover all parts thoroughly.
2. Apply in Early Spring, just before buds break.

### DORMANT COMBINATION SPRAY (for use in Northwest only)

MILLER'S SPRA-OIL	MILLER'S POLYGUL	Water
5 tablespoons	8 tablespoons	1 gallon
1 cup	3/4 pint	3 gallons
1 quart	1 1/2 quarts	12 1/2 gallons

NET CONTENTS  
ONE QUART



# Spra-oil

FOR CONTROL  
OF  
**RED SPIDER MITES, MEALY BUGS  
CRAWLER SCALE, HOLLY SCALE  
SOFT BROWN SCALE**

ACTIVE INGREDIENT                      By wt.  
Petroleum Oil                              98%  
INERT INGREDIENTS                      2%  
Mix. Unlabeled Residue 92%  
Grade: Unlabeled

**CAUTION** Keep out of the reach of children  
See right side panel for additional cautions

E.P.A. Reg. No. 802-179-AA

# Miller's SPRA-OIL

PLEASE READ ENTIRE LABEL USE ONLY AS DIRECTED,  
NOTING ALL CAUTIONS AND WARNINGS.

## CAUTION

Harmful if swallowed. If swallowed **do not** induce vomiting. Call a physician immediately.

Never apply a combination oil and sulphur spray on foliage. Do not apply an oil spray to foliage until 30 to 60 days have elapsed after a sulphur application as injury to foliage or flowers will result. Do not spray when weather is excessively hot. Be sure roots of plant are well watered before spraying.

Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Destroy empty container. Do not reuse for any purpose.

**NOTICE:** Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

MP 11 J

THE CHAS. H. LILLY CO.  
PORTLAND, OREGON  
EPA Est. 802-OR-1