

DIRECTIONS FOR USE — LIVESTOCK

LIVESTOCK SPRAYS AND DIPS

BEEF CATTLE — to control Lice and Ticks

- 1 Use at the rate of 1 1/4 pt per 100 gallons of water
- 2 Spray or dip to assure thorough and complete coverage

BEEF CATTLE — to control Mange Mites (Scab or Sarcoptic Mange)

- 1 Use at the rate of 1 3/4 pt per 100 gallons of water.
- 2 Complete, thorough coverage of entire animal is essential to obtain the best results
- 3 Make two applications only, one or two weeks apart

SHEEP AND GOATS — to control Lice, Ticks and Scab Mites (scabies)



- 1 **As a spray** - use 1 1/4 pt per 100 gallons of water
- 2 Apply to thoroughly wet animals
- 3 For Scab control, make 2 applications 10 days apart
- 4 **As a dip** — use 1 1/4 pt per 100 gallons of water. For Scab control, make 2 dippings 10 days apart.
- 5 Do not use on dairy goats

SWINE — to control Lice and Sarcoptic Mange

- 1 Use 1 3/4 pt per 100 gallons of water
- 2 Spray thoroughly but do not over wet

USE PRECAUTIONS — LIVESTOCK

- 1 Use only specified amount of chemical when mixing spray or dip. Make sure there is thorough and complete mixing by proper agitation before using spray or dip solution.
- 2 Do not spray within 30 days of slaughter. Do not dip within 60 days of slaughter.
- 3 Do not treat sick or emaciated animals.
- 4 Do not spray or dip animals under 3 months of age.
- 5 When necessary to spray young animals (over 3 months of age), use 1 pt per 100 quarts of water.
- 6 Do not treat sows within 2 weeks of farrowing nor for at least 3 weeks thereafter.
- 7 Do not use in poultry houses. Do not use on dairy cattle, or on dairy goats.
- 8 **CAUTION** — as a general precaution in spraying or dipping animals, avoid treating during cold, stormy weather. Water animals before dipping to prevent them from drinking the dip.



LINDANE 20E

(Contains 1.75 lbs. Lindane per gallon)

LIVESTOCK...FLY CONTROL...ORNAMENTALS

ACTIVE INGREDIENT

Lindane (Gamma Isomer of Benzene

Hexachloride) 20%

INERT INGREDIENTS 80%

E.P.A. Reg. No 802-186-ZA

PROTECT FROM UNDUE FREEZING TEMPERATURES

CAUTION — Keep out of reach of children

See right side panel for additional cautions

NET CONTENTS: _____ GALLON(S)

THE CHAS. H. LILLY CO.
PORTLAND, OREGON
EPA Est. 802-OR-1



Miller's



LINDANE 20E

(Contains 1.75 lbs. Lindane per gallon)

LIVESTOCK...FLY CONTROL...ORNAMENTALS

ACTIVE INGREDIENT

Lindane (Gamma Isomer of Benzene Hexachloride)	20%
INERT INGREDIENTS	80%

E.P.A. Reg. No. 802-186-ZA

PROTECT FROM UNDUE FREEZING TEMPERATURES

CAUTION — Keep out of reach of children

See right side panel for additional cautions

NET CONTENTS: _____ GALLON(S)

THE CHAS. H. LILLY CO.
PORTLAND, OREGON
EPA Est. 802-OR-1



Miller's

FLY CONTROL (House and Stable Flies)

- Residual Spray for Barns and Outbuildings**—use 1 1/2 gals. per 100 gal. water. Apply as a coarse spray to thoroughly wet walls, windows, fences and exterior surfaces to point of runoff.
- Interior Spray for Barns, Outbuildings**—use 1 1/2 gals. per 100 gals. of water. Apply as a coarse spray to thoroughly wet walls, ceilings, beams, screens and floor using approximately 1 gal. diluted spray per 1,000 sq. ft.
- Do not use in dairy barns or milk rooms. Do not contaminate animals' feed or water.

SYMPHYLID CONTROL ON ORNAMENTALS

Use at the rate of 1/2 pt. per 1,000 sq. ft. and work thoroughly into the top six inches of soil. This will also aid in the control of root weevil larvae and white grubs. Do not use on soil to be used for production of root or vegetable crops as Lindane may cause an off-flavor.

CAUTION

Harmful if swallowed. May be absorbed through skin. Avoid breathing spray mist. Avoid contact with skin and eyes. In case of contact, wash immediately with soap and water. Wash hands thoroughly with soap and water after handling the concentrate or applying the diluted product to animals.

Avoid contamination of feed and foodstuffs. Do not use on household pets or humans. This product is toxic to fish, birds and other wildlife. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes.

DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME.

NOTICE: Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given hereon.

M.P.C.

DIRECTIONS FOR USE -- LIVESTOCK

LIVESTOCK SPRAYS AND DIPS

BEEF CATTLE -- to control Lice and Ticks.
1. Use at the rate of 1 1/4 pt. per 100 gallons of water.
2. Spray or dip to assure thorough and complete coverage.

BEEF CATTLE -- to control Mange Mites (Scab) or Sarcoptic Mange.
1. Use at the rate of 1 3/4 pt. per 100 gallons of water.

2. Complete, thorough coverage of entire animal is essential to obtain the best results.
3. Make two applications only, one or two weeks apart.

SHEEP AND GOATS -- to control Lice, Ticks and Scab Mites (scabies).

1. As a spray - use 1 1/4 pt. per 100 gallons of water.
2. Apply to thoroughly wet animals.
3. For Scab control, make 2 applications 10 days apart.
4. As a dip -- use 1 1/4 pt. per 100 gallons of water. For Scab control, make 2 dippings 10 days apart.
5. Do not use on dairy goats.

SWINE -- to control Lice and Sarcoptic Mange.

1. Use 1 3/4 pt. per 100 gallons of water.
2. Spray thoroughly but do not over wet.

USE PRECAUTIONS -- LIVESTOCK

1. Use only specified amount of chemical when mixing spray or dip. Make sure there is thorough and complete mixing by proper agitation before using spray or dip solution.
2. Do not spray within 30 days of slaughter. Do not dip within 60 days of slaughter.
3. Do not treat sick or emaciated animals.
4. Do not spray or dip animals under 3 months of age.
5. When necessary to spray young animals (over 3 months of age), use 1 pt. per 100 gals of water.
6. Do not treat sows within 2 weeks of farrowing nor for at least 3 weeks thereafter.
7. Do not use in poultry houses. Do not use on dairy cattle, or on dairy goats.
8. CAUTION -- as a general precaution in spraying or dipping animals, avoid treating during cold, stormy weather. Water animals before dipping to prevent them from drinking the dip.

802-186



LIVESTOCK



Microptic Mangel

is essential to obtain the best

Mites (scabies)

For Scab control, make 2

LIVESTOCK

spray or dip. Make sure there
before using spray or dip solu-

within 60 days of slaughter

of age), use 1 pt per 100

at least 3 weeks thereafer
cattle, or on dairy goats

feeding animals, avoid treat
feeding to prevent them from



LINDANE 20E

(Contains 1.75 lbs. Lindane per gallon)

LIVESTOCK...FLY CONTROL...ORNAMENTALS

ACTIVE INGREDIENT

Lindane (Gamma Isomer of Benzene
Hexachloride)

INERT INGREDIENTS

ACCEPTED
802-186
20%
80%
27

E.P.A. Reg. No. 802-186-2A

PROTECT FROM UNDUE FREEZING TEMPERATURES

CAUTION — Keep out of reach of children

See right side panel for additional cautions

NET CONTENTS: _____ GALLON(S)

THE CHAS. H. LILLY CO.
 PORTLAND, OREGON
 EPA Est. 802-OR-1




RECEIVED
 5 NOV 1974
 REGISTRATION DIVISION
 PESTICIDES, EPA

Miller's

LINDANE 20E

(5 lbs. Lindane per gallon)

CONTROL...ORNAMENTALS

Derivative of Benzene

Reg. No 802-186-ZA

AVOID FREEZING TEMPERATURES

Keep out of reach of children

See panel for additional cautions

TS: _____ GALLON(S)

LILLY CO.
OREGON
972-OR-1

ACCEPTED
802-186
20%
80%
27 1976

Miller

FLY CONTROL (House and Stable Flies)

- Residual Spray for Barns and Outbuildings**—use 1 1/2 gals. per 100 gals. of water. Apply as a coarse spray to thoroughly wet walls, windows, fences and other exterior surfaces to point of runoff
- Interior Spray for Barns, Outbuildings**—use 1 1/2 gals. per 100 gals. of water. Apply as a coarse spray to thoroughly wet walls, ceilings, beams, screens and floors, using approximately 1 gal. diluted spray per 1,000 sq. ft.
- Do not use in dairy barns or milk rooms. Do not contaminate animals' feed or water.

SYMPHYLID CONTROL ON ORNAMENTALS

Use at the rate of 1/2 pt. per 1,000 sq. ft. and work thoroughly into the top six inches of soil. This will also aid in the control of root weevil larvae and white grubs. Do not use on soil to be used for production of root or vegetable crops as Lindane may cause an off-flavor.

CAUTION

Harmful if swallowed. May be absorbed through skin. Avoid breathing spray mist. Avoid contact with skin and eyes. In case of contact, wash immediately with soap and water. Wash hands thoroughly with soap and water after handling the concentrate or applying the diluted product to animals.

Avoid contamination of feed and foodstuffs. Do not use on household pets or humans.

This product is toxic to fish, birds and other wildlife. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes.

DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME.

NOTICE: Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

802-186

5