

DIRECTIONS

APPLES and PEARS


1. **Apples** . . . use 2 level teaspoons per quart of water (1/2 cup per 3 gallons).
2. **Pears** . . . use 1 1/2 level teaspoons per quart of water (6 level tablespoons per 3 gallons).
3. Make **first** application before opening of any blossoms (Pre-Pink Spray); **second**, just before blossoms open (Pink Spray); **third**, when most of the petals have fallen (Calyx Spray). Do not apply within 21 days of harvest.
4. For control of Powdery Mildew later in the season, consult your State Agricultural Experiment Station or your State Extension Service Authorities for material to use as recommendations vary in different localities and on various varieties.

GREENHOUSE Roses, African Violets, Asters, Larkspur, Begonias, Belladonna, Dahlias, Chrysanthemums, Calendula, Transvaal Daisies, Hydrangeas, Snapdragons

1. Use 3/4 level teaspoon per quart of water (3 tablespoons per 3 gallons).
2. Make first application as a protective spray when mildew may be prevalent and repeat at 4 to 5 day intervals until mildew is under control.
3. Apply as a fine mist spray.
4. Be sure underside of leaves is covered.
5. Apply with care when temperatures in greenhouse approach 85° and do not apply when it goes over 90°.


CAUTION . . . avoid prolonged skin contact. May cause skin irritation or sensitization. Wash thoroughly. Keep out of eyes. Avoid inhalation of mist or dust. Harmful if swallowed. Avoid contamination of foodstuffs. Wash the edible portions of fruits before using. Do not use with oil or oil base sprays or with liquid lime sulphur. Destroy empty container. Do not re-use for any purpose.

NOTICE: . . . We warrant the careful following and method of application as stated on the label. We do not warrant the effectiveness of the product as recommended on the label. The use of this product is at the user's risk. We are not responsible for any damage or loss of crops or other property caused by the use of this product.



KARASpra

Karathane® Spray Powder



ROSES — FLOWERS SHRUBS — TREES — FRUITS NURSERY & GREENHOUSE

Active ingredients:
 *2,4-Dinitro-6-octyl phenyl crotonate 3.17%
 *2,6-Dinitro-4-octyl phenyl crotonate 0.21%
 -Nitrooctyl phenols — principally dinitro 93.62%
 Inert ingredients
 *A mixture of 1-methylheptyl, 1-ethylhexyl, 1-propylpentyl isomers

CAUTION — Keep out of reach of children

See left side panel for additional cautions.

E P A Reg No 802-193-AA

Net weight 1 lb.

MILLER PRODUCTS

PORTLAND, OREGON 97218

DIRECTIONS

GENERAL DIRECTIONS . . . In the control of Powdery Mildew, Miller's KARASpra is most effective when used in a protective spray schedule at any time during mildew (moist, warm) weather. Repeat sprays at 4 to 5 day intervals until mildew is under control. Do not apply outdoors when temperature will exceed 90-95° F. during day.

ROSES . . . Outdoors, including the Rosa Multiflora

1. Use level teaspoon per quart of water (4 level tablespoons per 3 gallons).
2. If Black Spot is also a problem, add Miller's Garden CAPTAN to the spray.

FLOWERS . . . Outdoors . . . Begonias, Asters, Larkspur, Belladonna, Dahlias, Chrysanthemums, Calendula, Transvaal Daisies, Hydrangeas, Snapdragons

1. Use 1 level teaspoon per quart of water (4 level tablespoons per 3 gallons).

ORNAMENTAL TREES and SHRUBS . . . Hydrangeas, Euonymus

1. Use 1 1/2 level teaspoons per quart of water (6 level tablespoons per 3 gallons).

NURSERY STOCK . . . Apples, Peaches, Cherries (non-bearing)

1. Use 1 1/2 level teaspoons per quart of water (6 level tablespoons per 3 gallons).
2. Make first application as buds begin to break and repeat sprays at 4 to 5 day intervals until mildew is under control.