

*Miller's* **SULFO 95**  
**WETTABLE SULPHUR**

E.P.A. REG. No. 802- 297-1A

**DIRECTIONS FOR USE**

**PEACHES**—Powdery Mildew  
**Shuck Fall Spray**—use 4 to 5 lbs. per 100 gal. water (400 to 600 gal. per acre)  
**2 to 3 weeks after Shuck Fall Spray**—use 3 lbs. per 100 gal. of water (600 to 800 gal. per acre)

**PEACHES**—Brown Rot  
**Blossom, Petal Fall, Shuck Fall and 2 weeks later**—use 5 lbs. per 100 gal. of water (400 to 600 gal. per acre)  
**Pre-harvest and during harvest**—use 5 lbs. per 100 gal. water (600 to 800 gal. per acre), applying at 20 and 10 days before harvest and at picking time

**CHERRIES**—Powdery Mildew  
**Shuck Fall and 2 to 3 weeks later**—use 5 lbs. per 100 gal. of water (apply 400 to 600 gal. per acre for shuck fall, 600 to 800 gal. per acre later)  
**Post-Harvest**—use 5 lbs. per 100 gal. of water (600 to 800 gal. per acre). This spray may be applied to reduce infection

**CHERRIES**—Brown Rot  
**Blossom, Petal Fall, Shuck Fall and Summer Sprays**—use 5 lbs. per 100 gal. of water (400 to 600 gal. per acre)  
**Pre-harvest and during harvest**—use 5 lbs. per 100 gal. water (600 to 800 gal. per acre), applying at 20 and 10 days before harvest and at picking time

**PRUNES and PLUMS**—Brown Rot  
**Blossom, Petal Fall, Shuck Fall and Summer Sprays**—use 5 lbs. per 100 gal. water (400 to 600 gal. per acre)

**GRAPES**—Powdery Mildew  
**During growing season**—use 2 lbs. per 100 gal. water (300 to 400 gal. per acre). Make applications at following seasons: (1) When the new shoot growth is 6 to 10 inches long. (2) Just before blossoming. (3)

**DIRECTIONS FOR USE CONTINUED**

After the berries are set and not later than one fourth grown. (4) Between the time the berries are one-half and fully grown. (5) Just as the grapes begin to ripen

**ROSES, PERENNIALS, SHRUBS**—Black Spot, Powdery Mildew, Rust  
 Use 6 lbs. per 100 gal. of water and spray for full coverage, wetting both sides of leaves. Begin applications when new growth appears in the spring and repeat at 4 to 7 day intervals as necessary. **CAUTION**—some burning may occur when used during warm weather.

**MIXING DIRECTIONS**

Pour recommended amount of this material through screen on surface of water in nearly filled spray tank, or pre-mix in a bucket half filled with water and pour mixture into nearly filled spray tank. Add balance of water to fill tank. Keep agitator running during filling and spraying operations. Do not allow mixture to stand. Do not use in low volume gear-type equipment. Do not combine with emulsifiable liquids in the same spray tank.

**CAUTION — Keep out of reach of children**

Do not apply when shade temperatures exceed or are likely to exceed 90° F. Sulphur may burn foliage and fruit during periods of high temperatures and under certain climatic conditions. Do not use sulphur within 30 days of an oil spray. Do not reuse container, do not burn or empty.  
 If the crop to be treated will be used for canning or freezing purposes, consult food processor before applying sulphur.  
 Consult State Agricultural Experiment Station or State Extension Service Authorities for exact timing and spacing of sprays in your area.

Active ingredients	
Sulphur	95.0%
Inert ingredients	5.0%

**NOTICE:** Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not applicable. Do not apply and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material but in strict accordance with directions given hereon.

MP 111

Manufactured for ...  
 THE CHAS. H. LILLY CO.  
 PORTLAND, OREGON

NET WT. **50 LBS.**