

Insist on ...

Miller's



# CHLORDANE 20G

(20% Chlordane Granules)

**SOIL INSECTICIDE**

**FOR VEGETABLES—BERRIES—TURF—SHRUBS—NURSERIES**

## DIRECTIONS FOR USE

### VEGETABLES

**BEANS, BEETS, COLE CROPS (Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Collards, Kale, Kohlrabi), CANTALOUPE, CELERY, CUCUMBERS, EGGPLANTS, LETTUCE, MELONS, OKRA, ONIONS, PEAS, PEPPERS, PIMENTOS, POTATOES, RADISHES, RUTABAGAS, SQUASH (Summer and Winter), SWEET POTATOES, TURNIPS, TOMATOES**

**Cutworms, Wireworms, White Grubs, Flea Beetle Larvae:**

Apply 20 lbs. per acre as a preplanting soil treatment. Thoroughly incorporate into top 3 to 6 inches of soil before planting.

**Asiatic Garden Beetle Larvae, White Fringed Beetle Larvae:**

Apply 25 lbs. per acre as a preplanting soil treatment. Mix thoroughly into top 3 to 6 inches of soil before planting.

**Japanese Beetle Larvae, Wireworms:**

Apply 40 to 50 lbs. per acre as a preplanting soil treatment. Thoroughly mix into top 3 to 6 inches of soil before planting.

### BERRIES

**BLACKBERRY BLUEBERRY BOYSENBERRY DEWBERRY  
HUCKLEBERRY LOGANBERRY RASPBERRY STRAWBERRY  
YOUNGBERRY**

**Strawberry Root Weevil Larvae:** Apply 20 lbs. per acre as a preplanting soil treatment.

## CAUTION—Keep out of reach of children

Avoid inhalation of dust. Avoid contact with skin, eyes and clothing. Wash thoroughly with soap and water after handling.

Avoid contamination of feed and feedstuffs.

## USE PRECAUTIONS

Do not contaminate drinking water or use near fishponds.

Keep children and pets off treated turf and lawns until the material has been thoroughly washed into the soil by watering or by rain and the grass has completely dried.

Do not graze livestock on treated areas nor feed clippings from treated areas to livestock.

This product is highly toxic to bees exposed to direct treatment or residue on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

This product is toxic to fish, birds and other wildlife. Keep out of lakes, streams and ponds. Birds and other wildlife in treated areas may be killed. Do not apply when weather conditions favor drift from areas treated. Do not apply when runoff is likely to occur. Do not use in or near water bodies or in areas of environmental sensitivity. Avoid

MELONS, OKRA, ONIONS, PEAS, PEPPERS, PIMENTOS, POTATOES, RADISHES, RUTABAGAS, SQUASH (Summer and Winter), SWEET POTATOES, TURNIPS, TOMATOES

**Cutworms, Wireworms, White Grubs, Flea Beetle Larvae:** Apply 20 lbs. per acre as a preplanting soil treatment. Thoroughly incorporate into top 3 to 6 inches of soil before planting.

**Asiatic Garden Beetle Larvae, White Fringed Beetle Larvae:** Apply 25 lbs. per acre as a preplanting soil treatment. Mix thoroughly into top 3 to 6 inches of soil before planting.

**Japanese Beetle Larvae, Wireworm:** Apply 40 to 50 lbs. per acre as a preplanting soil treatment. Thoroughly mix into top 3 to 6 inches of soil before planting.

**BERRIES**

**BLACKBERRY, BLUEBERRY, BOYSENBERRY, DEWBERRY, HUCKLEBERRY, LOGANBERRY, RASPBERRY, STRAWBERRY, YOUNGBERRY**

**Strawberry Root Weevil Larvae:** Apply 40 to 50 lbs. per acre. **Preplanting Soil Treatment:** Thoroughly mix into top 6 inches of soil before setting out plants. **Post-Planting Soil Treatment:** Make application at base of plants and incorporate into soil. Do not apply when fruit is present.

**LAWNS AND TURFS—Sod Webworm:** Apply evenly over the turf at the rate of 15 lbs. per acre. Water treated area thoroughly after application. Early spring applications will give best results. Keep children and pets off treated lawns or turf until the material has been thoroughly washed into the soil by watering or by rain and the grass has completely dried.

**NURSERIES, SHRUB BEDS—Strawberry Root Weevil Larvae, Cutworms, Black Vine Weevil Larvae:** Apply 40 lbs. per acre (1 lb. per 1000 sq. ft.) as a preplanting soil treatment. Work thoroughly into top 4 to 6 inches of soil before planting.

**USE PRECAUTIONS**

Do not contaminate drinking water, or use near fishponds.

Keep children and pets off treated turf and lawns until the material has been thoroughly washed into the soil by watering or by rain and the grass has completely dried.

Do not graze livestock on treated areas nor feed clippings from treated areas to livestock.

This product is highly toxic to bees exposed to direct treatment or residue on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

This product is toxic to fish, birds, and other wildlife. Keep out of lakes, streams and ponds. Birds and other wildlife in treated areas may be killed. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes.

**ACTIVE INGREDIENT**

\*Chlordane, Technical . . . . . 20.0%

**INERT INGREDIENTS** . . . . . 80.0%

\*Equivalent to 12% octachloro-4,7-methanotetrahydroindane and 8% related compounds.

**E.P.A. Reg. No. 802-234-AA**

**NOTICE:** Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage and handling of this material not in strict accordance with directions given herewith. MP 6K

**ACCEPTED**  
DEC 10 1975  
Under the Federal Insecticide, Fungicide and Rodenticide Act, registered for the pesticide

**NET WT. \_\_\_\_\_ LBS.**

**THE CHAS. H. LILLY CO.**  
**PORTLAND, OREGON 97218**  
EPA Est 802 OR 1

