

ACCEPTED

MAY 29 1975

Under the Federal Insect Act, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 802-235

Miller's CAPTAN 5D

(5% Captan Dust)
AGRICULTURAL FUNGICIDE

ACTIVE INGREDIENT

Captan (N-[(trichloromethyl) thio]-4-cyclohexene-1,2-dicarboximide) 5.0%

INERT INGREDIENTS 95.0%

E.P.A. Reg. No. 802-235-AA

CAUTION - Keep out of reach of children

Avoid inhalation of dust. Avoid contact with skin. In case of contact, wash with soap and water. Avoid contamination of feed and foodstuffs.

This product is toxic to fish. Keep out of lakes, ponds and streams. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Consult State Agricultural Experiment Station or State Extension Service Authorities as to the correct timing and number of applications necessary for your area consistent with label directions and cautions.

Dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes.

GENERAL DIRECTIONS

Apply on first appearance of diseases. Dust to thoroughly cover all parts of plants. Repeat applications at 4 to 7 day intervals as needed to maintain control.

SUGGESTED USES

Crop and Pest Lbs. per Acre

STRAWBERRIES - Fruit Rot (Botrytis cinerea) 50

Make application as soon as possible after each rainy period, but no more than 2 during each 14 day period. During wet bloom period make 3 to 4 applications; 2 to 3 applications if bloom period is dry. No application during harvest is required.

Crop and Pest Lbs. per Acre

APRICOTS, PEACHES - Brown Rot 60 to 85
Apricots - apply at red bud stage and repeat at 85% petal fall. Peaches - apply in the blossom and pre-harvest applications.

CUCURBITS (Cantaloupe, Cucumbers, Watermelons, Summer and Winter Squash) - Angular Leaf Spot, Anthracnose, Downy Mildew 40 to 60*
*on young plants, increasing to 60 to 75 lbs. per acre on mature plants. Start applications when runners begin to form or at first sign of disease.

TOMATOES - Early and Late Blight, Anthracnose Grey Leaf Spot, Septoria Leaf Spot 40 to 60*
*on young plants; 65 to 85 lbs. per acre on mature plants. Make weekly applications.

POTATOES - Early and Late Blight 60 to 70*
*on young plants; 80 lbs. per acre on mature plants. Make first application when plants are 2 to 4 inches high. Repeat at 7 to 10 day intervals as long as disease threatens.

POTATOES - Seed Piece Treatment -- Damping Off, Seed Rot, Brown Eye Disease, Verticillium Wilt.

Use at rate of 20 oz. per bushel of cut seed pieces, or 32 oz. (2 lbs.) to 100 lbs. of cut seed pieces. Apply in convenient container or by duster attachment over belt. Coat each seed piece evenly with a light film of dust. Treat pieces within 6 hours of cutting. If the planting is delayed, store treated seed pieces in open crates, where there is a good air movement, for 2 days before bagging. Do not use treated seed for food or feed purposes. Mark bags TREATED SEED PIECES.

NOTICE: Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material, not in strict accordance with directions given hereon.

MP-5K

NET WT. 50 LB

THE CHAS. H. LILLY CO.
PORTLAND, OREGON
EPA Est. 802-OR-1

