

Miller's CAPTAN 5D

(5% CAPTAN DUST -- AGRICULTURAL FUNGICIDE)

ACCEPTED
 APR 7 1979
 Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 802-235.

ACTIVE INGREDIENTS: Captan (N-[(trichloromethyl) thio]-4-cyclohexene-1,2-dicarboximide) 5.0%
INERT INGREDIENTS: 95.0%
 E.P.A. Reg. No. 802-235

GENERAL DIRECTIONS:
 Apply on first appearance of diseases. Dust to thoroughly cover all parts of plants. Repeat applications at 4 to 7 day intervals as needed to maintain control.

Keep out of reach of children
CAUTION

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION: Avoid inhalation of dust. Avoid contact with skin. Avoid contamination of feed and foodstuffs.
STATEMENT OF PRACTICAL TREATMENT
 In case of contact, wash with soap and water.
ENVIRONMENTAL HAZARD
 This product is toxic to fish. Keep out of lakes, ponds and streams. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL
 Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Pesticide, spray mixture, or rinse that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies.
Container disposal: Dispose of in an incinerator or landfill approved for pesticide containers, or bury in a safe place.
 Consult federal, state or local disposal authorities for approved alternative procedures such as limited open burning.

Crop and Pest	Lbs. per Acre
STRAWBERRIES – Fruit Rot (<i>Botrytis cinerea</i>) Make application as soon as possible after each rainy period, but no more than 2 during each 14 day period. During wet bloom period make 3 to 4 applications; 2 to 3 applications if bloom period is dry. No application during harvest is required.	50
APRICOTS, PEACHES – Brown Rot Apricots -- apply at red bud stage and repeat at 85% petal fall. Peaches -- apply in the blossom and pre-harvest applications.	60 to 85
CUCURBITS (Cantaloupe, Cucumbers, Watermelons, Summer and Winter Squash) – Angular Leaf Spot, Anthracnose, Downy Mildew *on young plants, increasing to 60 to 75 lbs. per acre on mature plants. Start applications when runners begin to form or at first sign of disease.	40 to 60*
TOMATOES – Early and Late Blight, Anthracnose, Grey Leaf Spot, Septoria Leaf Spot *on young plants; 65 to 85 lbs. per acre on mature plants. Make weekly applications.	40 to 60*
POTATOES – Early and Late Blight *on young plants; 80 lbs. per acre on mature plants. Make first application when plants are 2 to 4 inches high. Repeat at 7 to 10 day intervals as long as disease threatens.	60 to 70*

NOTICE: Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this product not in strict accordance with directions given herewith.
 LC- 60

NET WT. 50 LBS.

THE CHAS. H. LILLY Co.
 PORTLAND, OREGON 97218
 EPA Est. 802-OR-1



ACTIVE INGREDIENTS:

Captan (N-[(trichloromethyl)thio]-4-cyclohexene-1,2-dicarboximide) 5.0%

INERT INGREDIENTS: 95.0%

EPA Reg. No. 802-235

Keep out of reach of children

CAUTION

PRECAUTIONARY STATEMENTS

HAZARD TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Avoid inhalation of dust. Avoid contact with skin. Avoid contamination of feed and foodstuffs.

STATEMENT OF PRACTICAL TREATMENT

In case of contact, wash with soap and water.

ENVIRONMENTAL HAZARD

This product is toxic to fish. Keep out of lakes, ponds and streams. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

NOTICE: Follow directions carefully. Timing and method of application and crop conditions, mixture with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material. It is strict accordance with directions given herewith.

LC-10

NET WT. 50 LBS.

**THE CHAS. H. LILLY CO.
PORTLAND, OREGON 97218**

EPA Est. 802-OR-1

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

Pesticide, spray mixture, or rinsate that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies.

Container disposal: Dispose of in an incinerator or landfill approved for pesticide containers, or bury in a safe place.

Consult federal, state or local disposal authorities for approved alternative procedures such as limited open burning.

GENERAL DIRECTIONS

POTATOES - SEED PIECE TREATMENT

Damping Off, Seed Rot, Brown Eye Disease, Verticillium Wilt

Use at rate of 20 oz. per bushel of cut seed pieces, or 32 oz. (2 lbs.) to 100 lbs. of cut seed pieces.

Apply in convenient container or by duster attachment over belt. Coat each seed piece evenly with a light film of dust.

Treat pieces within 6 hours of cutting. If the planting is delayed, store treated seed pieces in open crates, where there is good air movement, for 2 days before bagging.

DO NOT USE TREATED SEED FOR POTATOES.

Mark bags TREATED SEED PIECE

ACCEPTED
APR 7 1978
KERR
Under the Federal Insecticide, Fungicide and Rodenticide Act, registered for the pesticide. Registered under EPA Reg. No. 802-235

LILLY/MILLER