or preharvest ap-

t as soon as corn when corn is over

pray off foliage of

spray off foliage

reatment calculate

f actively growing

20 to 35 lbs.

fter application and

of water and apply

rer heads; {2} when

elly temporary. This Buffalo, Dichondra



# ENVY 2,4-D

(4 lbs. 2,4-D acid equivalent per gallon) Solubilized 2,4-D Acid A special formulation for control of . . .

Russian Knapwood Wild Garlic **Leafy Spurge** 

**Canada Thistle** 

**Blue Flowering** Letture **Morning Glory** 

Wiid Onlon St. Johnswort White Top **Tansy Ragwort** 

on Fallow Land. Fence Rows. Roadsides. Non-crop Land

ACTIVE INGREDIENTS	
Triethylamine Salt of 2,4-Dichlorophenoxyacetic Acid*	50.32
2,4-Dichlorophenoxyacetic Acid	10.00
INERT INGREDIENTS	39.68
*2,4-Dichlorophenoxyacetic Acid equivalent 34.5%. Total 2,4-Dichloropheno	XYace
Acid equiv. and Actual 44.5%.	

E.P.A. Reg. No. 802-241-ZA


WARNING - FLAMMABLE! KEEP AWAY FROM HEAT OR OPEN FLAME.

CAUTION Keep out of reach of children
See right side panel for additional cautions.

**NET CONTENTS** 

GALLONS

THE CHAS. H. LILLY CO. PORTLAND, OREGON EPA Est. 802-OR-1



## **GRASS SEED CROPS**

Use 1 to 1 1/2 pts, per acre. New seedlings in the spring should be sprayed after the grasses have at least five leaves. Where irrigation is practiced regularly, application may be made after the two leaf stage. When small seedlings are sprayed do not use more than 1 pt. per acre.

FALL SEEDED STANDS IN WESTERN OREGON - Use 1 pt. per acre any time after 2 leaf stage.

ESTABLISHED STANDS — Use 1 to 1 1/2 pts. per acre. Application may be made in the fall or early spring, prior to the time seed heads appear in the boot.

# CAUTION

### Please read entire label. Use only as directed, noting all cautions and warnings

### Harmful if swallowed.

This product may cause skin irritation. Avoid contact with skin, eyes and clothing. In case of contact, wash skin with plenty of soap and water; flush eyes with pienty of water for at least 15 minutes and get medical attention. Keep out of reach of children and domestic animals.

Do not store with foodstuffs. Do not transport or store near fertilizers, insecticides, fungicides or seeds.

Do, not apply directly to or otherwise permit sprays to come in contact with vegetables, flowers, grapes, fruit trees, cane berries, ornamentals or other desirable plants which are sensitive to 2,4-D and do not permit spray mist to drift onto them, since even minute quantities of the spray may cause severe injury during both growing and dormant periods. Coarse sprays are less likely to drift.

Application by airplane, ground equipment or hand sprayers should be made only when there is no hazard from drift.

Do not apply by airplane in the vicinity of grapes or other desirable 2,4-D susceptible plants.

Do not contaminate irrigation ditches or water used for domestic purposes. Excessive amounts of 2,4-D in the soil may temporarily inhibit seed germination or plant growth.

At higher temperatures, vaporization may cause injury to susceptible plants growing nearby. Keep container tightly closed when not in use.

Do not use spray equipment for any other purpose. Such equipment should not be used for application of insecticides, fungicides, or other agricultural chemicals.

Be sure that use of this product conforms to all applicable local regulations.

Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Do not contaminate water by cleaning of equipment or disposal of wastes. Containers should be disposed of by punching holes in them, crushing and burying with wastes.

NOTICE: Follow directions carefully. Timing and method of application, weather and crop conditions, mixture, with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

MP-7K