



# ENVY 2,4-D

A special formulation for control of . . .

Russian Knapweed	Blue Flowering	Wild Onion
Wild Garlic	Lettuce	St. Johnswort
Leafy Spurge	Morning Glory	White Top
Canada Thistle		Taney Ragwort

on Fallow Land, Fence Rows, Roadides, Non-crop Land

**ACTIVE INGREDIENTS**  
 Triethylemine Salt of 2,4-Dichlorophenoxyacetic Acid\* . . . . . 46.85%  
 2,4-Dichlorophenoxyacetic Acid . . . . . 9.50%  
**INERT INGREDIENTS** . . . . . 41.85%  
 \*2,4-Dichlorophenoxyacetic Acid equivalent 33.36%, Total 2,4-Dichlorophenoxyacetic Acid equivalent and Actual 42.86% - 3.8 lbs./gal. T  
 † Tester specific by AOAC Method No. 8. D01.†

**WARNING - FLAMMABLE! KEEP AWAY FROM HEAT OR OPEN FLAME.**

**CAUTION** Keep out of reach of children.  
 See right side panel for additional cautions.

NET CONTENTS \_\_\_\_\_ GALLONS

THE CHAS. H. LILLY CO., PORTLAND, OREGON 97218 • EPA Est. 802-08-1 EPA Reg. No. 802-241

**GRASS SEED CROPS**

Use 1 to 1 1/2 pts. per acre. New seedings in the spring should be sprayed. Where irrigation is practiced regularly, application may be made after the seedlings are established. Do not use more than 1 pt. per acre.

**FALL SEEDED STANDS IN WESTERN OREGON** - Use 1 pt. per acre and  
**ESTABLISHED STANDS** - Use 1 to 1 1/2 pts. per acre. Application may be made at the time seed heads appear in the boot.

**CAUTION**

Please read entire label. Use only as directed, noting all directions. Store at temperatures above 41° F.

**Harmful if swallowed**

This product may cause skin irritation. Avoid contact with skin, eyes and clothing. In case of contact with skin, wash with plenty of water for at least 15 minutes. In case of contact with eyes, flush eyes with plenty of water for at least 15 minutes. Keep out of reach of children and domestic animals.

Do not store with foodstuffs. Do not transport or store near tentacles, etc.

Do not apply directly to or otherwise permit sprays to come in contact with cane berries, ornamentals or other desirable plants which are sensitive to 2,4-D. To them, since even minute quantities of the spray may cause severe injury. Coarse sprays are less likely to drift.

Application by airplane, ground equipment or hand sprayers should be made.

Do not apply by airplane in the vicinity of grapes or other desirable 2,4-D crops.

Do not contaminate irrigation ditches or water used for domestic purposes. This product may temporarily inhibit seed germination or plant growth.

At high temperatures, vaporization may cause injury to susceptible crops if closed when not in use.

Do not use spray equipment for any other purpose. Such equipment should be used only for fungicides or other agricultural chemicals.

Be sure that use of this product conforms to all applicable local regulations.

Wash spray equipment and containers and dispose of wastes by burying in soil. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not place in them or string and burying with wastes.

**NOTICE:** Follow directions carefully. Timing and method of application of other chemicals not specifically recommended and other influencing factors are the user's responsibility. Buyer assumes all risks of use, storage or handling of this product.

*Handwritten notes:*  
 10/1/81  
 10/1/81  
 10/1/81



# ENVY 2,4-D

A special formulation for control of...

<b>Russian Knapweed</b>	<b>Blue Flowering</b>	<b>Wild Onion</b>
<b>Wild Garlic</b>	<b>Lettuce</b>	<b>St. Johnswort</b>
<b>Leafy Spurge</b>	<b>Morning Glory</b>	<b>White Top</b>
<b>Canada Thistle</b>		<b>Tansy Ragwort</b>

on Fallow Land, Fence Rows, Roadside, Non-crop Land

**ACTIVE INGREDIENTS:**  
 Triethylamine Salt of 2,4-Dichlorophenoxyacetic Acid\* ..... 48.85%  
 2,4-Dichlorophenoxyacetic Acid ..... 8.50%  
**INERT INGREDIENTS** ..... 41.85%  
 \*2,4-Dichlorophenoxyacetic Acid equivalent 33.36% Total 2,4-Dichlorophenoxyacetic Acid equivalent and Actual 42.86% - 3.8 lbs/gal. 7  
 Testers specific by AOAC Method No. 6-DO1-5

**WARNING - FLAMMABLE! KEEP AWAY FROM HEAT OR OPEN FLAME.**

**CAUTION** Keep out of reach of children  
 See right side panel for additional cautions.

NET CONTENTS \_\_\_\_\_ GALLONS

THE CHAS. K. LILLY CO., PORTLAND, OREGON 97218 • EPA Est. 982-OR-1 EPA Reg. No. 982-241

### GRASS SEED CROPS

Use 1 to 1 1/2 pts. per acre. New seedlings in the spring should be sprayed after the grasses have at least five leaves. Where irrigation is practiced regularly, application may be made after the two leaf stage. When small seedlings are sprayed do not use more than 1 pt. per acre.

**FALL SEEDED STANDS IN WESTERN OREGON** - Use 1 pt. per acre any time after 2 leaf stage.

**ESTABLISHED STANDS** - Use 1 to 1 1/2 pts. per acre. Application may be made in the fall or early spring, prior to the time seed heads appear in the boot.

### CAUTION

Please read entire label. Use only as directed, noting all cautions and warnings.  
 Store at temperatures above 40° F.

Harmful if swallowed.

This product may cause skin irritation. Avoid contact with skin, eyes and clothing. In case of contact, wash skin with plenty of soap and water; flush eyes with plenty of water for at least 15 minutes and get medical attention. Keep out of reach of children and domestic animals.

Do not store with foodstuffs. Do not transport or store near fertilizers, insecticides, fungicides or seeds.

Do not apply directly to or otherwise permit sprays to come in contact with vegetables, flowers, grapes, fruit trees, cane berries, ornamentals or other desirable plants which are sensitive to 2,4-D and do not permit spray mist to drift on to them, since even minute quantities of the spray may cause severe injury during both growing and dormant periods. Coarse sprays are less likely to drift.

Application by airplane, ground equipment or hand sprayers should be made only when there is no hazard from drift.

Do not apply by airplane in the vicinity of grapes or other desirable 2,4-D susceptible plants.

Do not contaminate irrigation ditches or water used for domestic purposes. Excessive amounts of 2,4-D in the soil may temporarily inhibit seed germination or plant growth.

At higher temperatures, vaporization may cause injury to susceptible plants growing nearby. Keep container tightly closed when not in use.

Do not use spray equipment for any other purpose. Such equipment should not be used for application of insecticides, fungicides, or other agricultural chemicals.

Be sure that use of this product conforms to all applicable local regulations.

Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Do not contaminate water. Thorough cleaning of equipment or disposal of wastes. Containers should be disposed of by punching holes in them, crushing and burying with wastes.

**NOTICE:** Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herein.

140

EP 90