

Miller's **KELTHANE*** 4D

(4% Kelthane Dust. *Kelthane is the Reg. T.M. of the Rohm & Haas Co.)

AGRICULTURAL INSECTICIDE

ACTIVE INGREDIENT
 1,1-bis (chlorophenyl)-2,2,2-trichloroethanol 4.0%
 INERT INGREDIENTS 96.0%
 E.P.A. Reg. No. 802-251-AA

APPLICATION DIRECTIONS: Apply on first appearance of mites and repeat if necessary for adequate control.
 Do not apply closer to harvest than number of days indicated for the specific crop being treated.

CAUTION— Keep out of reach of children.

Avoid skin contact. Avoid inhalation of dust. Wash thoroughly after using.

Avoid contamination of feed and foodstuffs.


USE PRECAUTIONS

Read Entire Label. Use strictly in accordance with directions, cautions and warnings on label and in conformity with Federal and State Regulations.

Avoid drift to adjoining food and forage crops about to be harvested. Apply this product only as specified on this label. Do not apply when weather conditions favor drift of dust from areas treated. The grower is responsible for residues on his crops as well as for damages caused by drift from his property to that of others.

Toxic to fish. Keep out of lakes, streams and ponds. Do not apply where runoff is likely to occur.

Dispose of wastes by burying in non-crop lands away from water supplies. The containers should be disposed of by punching holes in them and burying with wastes. Do not contaminate water by cleaning of equipment, or disposal of wastes.



Crop and Pest	Pounds	Days per Acre
---------------	--------	---------------

ALFALFA-Seed Crops Only

Two-spotted and Pacific Mites	0	25
---	---	----

Use restricted to California, Idaho, Nevada, Oregon and Washington. Fields not to be grazed by livestock. Crop cuttings or residues not to be fed or baled. The fields must be cleared after harvest by burning or by discing into the soil.

BEANS - Dry, Green and Lima

Two-spotted and Pacific Mites	7	25 to 30
---	---	----------

Do not feed treated vines to meat or dairy animals.

CANTALOUPE, CUCUMBERS, MELONS, PUMPKINS, WINTER and SUMMER SQUASH, WATERMELONS

Two-spotted and Pacific Mites	2	25 to 30
---	---	----------

HOPS - Two-spotted Mites

Two-spotted Mites	7	25 to 35
-----------------------------	---	----------

TOMATOES, EGGPLANTS, PEPPERS - Two-Spotted,

Pacific and Tomato Russet Mites	2	25 to 35
---	---	----------

NOTICE: Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material, not in strict accordance with directions given herewith.

MP-5J

NET WT. 50 LBS.

THE CHAS. H. LILLY CO.
 PORTLAND, OREGON
 EPA Est. 802-OR-1

