

Miller's DITHANE* M22 6D

(6% Dithane M22 Dust)
AGRICULTURAL FUNGICIDE

Active ingredient	
Maneb (Manganese Ethylenebisdithiocarbamate) . . .	4.82
Inert ingredients	55.22

CAUTION—Keep out of reach of children

May cause irritation of nose, throat, eyes and skin. Avoid inhalation of dust. In case of skin contact, wash with water; for eyes wash copiously with water and get medical attention. Toxic to fish. Keep out of any body of water. Do not contaminate water by cleaning of equipment, or disposal of wastes.

Dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes.

Apply this product only as specified on this label. Avoid drift to adjoining food and forage crops about to be harvested. Do not apply when weather conditions favor drift of dust from areas treated. The grower is responsible for residues on his crops, as well as for damages caused by drift from his property to that of others.

STORAGE - Store in a dry place. Do not allow dust to become wet during storage. This may bring on chemical changes which will impair the fungicidal effectiveness of this dust and may also generate flammable vapors.

DIRECTIONS FOR USE

Use 17 to 45 lbs. per acre, depending upon size of plants.

Begin applications when diseases are reported in the area, or first appear, and continue at 7 to 10 day intervals, or as required for adequate control, unless otherwise specified for specific crop.

†Do not apply closer to harvest than number of days indicated for the specific crop being treated.

Crop and Pest Days

BEANS - Downy Mildew *

*Do not apply to string beans within 4 days of harvest.

CARROTS - Leaf Blight, Cercospora Leaf Spot. 0

CELERY - ^{Blight} Early and Late Blight 14

Begin dusting in the plant bed when the plants first come through the ground. Make further applications at 3 to 5 day intervals until plants are set in the field. Continue field applications on regular weekly schedule. Remove residue by stripping, trimming and washing.

CANTALOUPE, SUMMER SQUASH, WATERMELONS - Downy Mildew 5

CUCUMBERS - Anthracnose and Downy Mildew 5

On the above four crops, start applications when runners begin to form, or at first sign of disease. Repeat at weekly intervals. Direct dust to underside of leaves.

ONIONS - Downy Mildew, Purple Blotch 0

POTATOES - Early and Late Blight 0

SPINACH - Downy Mildew (Blue Mold) 10

Remove excess residues by washing.

TOMATOES - ^{Blight} Early and Late Blight, Anthracnose, Stemphylium (Gray Leaf Spot), Septoria Leaf Spots. 9

ACCEPTED
802-266
Nov. 14, 1973
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
ECONOMIC POISON REGISTER
ED UNDER NO. 802-266-266

Directions for use, storage, and disposal of containers, and other information, are given on the label. These should be read and followed in connection with the use of this product.

E.P.A. Reg. No. 802-266-AA

NET WT. 50 LB. BEST DOCUMENT AVAILABLE

THE CHAS. H. LILLY CO.
PORTLAND, OREGON

RECEIVED
OCT 12 1973
MILLERS

Miller's

Miller's DITHA

(6% Dithane
AGRICULTURA)

Active ingredient

Maneb (Manganese Ethylenedithiocarbamate)	4.8%
Inert ingredients	55.2%

CAUTION— Keep out of reach of children

May cause irritation of nose, throat, eyes and skin. Avoid inhalation of dust. In case of skin contact, wash with water; for eyes wash copiously with water and get medical attention.

Toxic to fish. Keep out of any body of water. Do not contaminate water by cleaning of equipment, or disposal of wastes.

Dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes.

Apply this product only as specified on this label. Avoid drift to adjoining food and forage crops about to be harvested. Do not apply when weather conditions favor drift of dust from areas treated. The grower is responsible for residues on his crops, as well as for damages caused by drift from his property to that of others.

STORAGE - Store in a dry place. Do not allow dust to become wet during storage. This may bring on chemical changes which will impair the fungicidal effectiveness of this dust and may also generate flammable vapors.

DIRECTIONS FOR USE

Use 17 to 45 lbs. per acre, depending upon size of plants.

Begin applications when diseases are reported in the area, or first appear, and continue at 7 to 10 day intervals, or as required for adequate control, unless otherwise specified for specific crop.

*Dithane is the Trade Mark of Rohm & Haas Co

E.P.A. Reg. No. 802-266-AA

NET WT. 50 LB.

THE CHAS.
PORTLAND