



Miller's SEVIN* 10D

(10% SEVIN* DUST)

REGISTERED TRADE MARK
LILLY & COMPANY

CAUTION — Keep out of reach of children

Do not breathe dust. Do not get dust on skin. Wash hands before eating. Take shower or bath after work. Wear goggles and long sleeved work clothing. Change to clean clothing daily.

Note to Physician: Carbofuran is a potent insecticide and irritant to the eyes. Avoid contact with eyes. If contact occurs, flush eyes with water and seek medical attention.

Do not use in areas where children play. Do not use in areas where water is used for drinking. Do not use in areas where water is used for washing dishes. Do not use in areas where water is used for washing of equipment.

Do not use in areas where people are exposed to direct sunlight. Do not use in areas where people are exposed to high temperatures. Do not use in areas where people are exposed to high humidity.

Crop and Pest	Lbs. per acre
APPLES Codling Moth, Green Apple Aphid, Apple Maggot, Red-headed Leaf Roller	10
PEACHES Peach Twig Borer, Oriental Fruit Moth	10
CHERRIES Cherry Fruit Fly	10
Repeat applications at 7 day intervals. CAUTION: Application of Carbofuran to apples within 50 days of full bloom may cause apple thinning. To avoid thinning effect, use alternative registered pesticide until 40 days after full bloom.	
STRAWBERRIES 1 Meadow Spittlebug, Strawberry Leaf-roller, Omnivorous Leaf Tier, Strawberry Fruit Worm	10
Apply when insects appear and repeat as needed.	
FILBERTS Filbert Worms	10
Apply on first appearance of adult moth and repeat 3 to 4 weeks later.	
Filbert Aphid and Filbert Leaf Roller	10
Time applications to treat when leaf roller eggs are hatching.	
Do not apply closer to harvest than specified on label for specific pest. Do not treat more than once for Filberts.	

On the following crops, apply at the rate specified on the label and repeat at 7 to 10 day intervals if needed. When used in accordance with label directions, the following crops may be produced for harvest for use as feed for dairy and meat animals and for use in processing plants and other uses.

Crop and Pest	Lbs. per acre
CORN Corn Earworm	10
Start dusting when silks first appear. Repeat at 7 to 10 day intervals throughout silking period. Direct dusting activity toward ear zone to insure thorough coverage of each ear. Application of this product to the tassel region of corn during the pollen shed period will seriously reduce pollen viability.	
POTATOES Colorado Potato Beetles, Leafhoppers, Flea Beetles	10
Dust thoroughly upper and lower leaf surfaces. Do not graze treated fields or feed vines to dairy or meat animals.	
BEANS Flea Beetles, Leafhoppers, Cucumber Beetles	10
Apply when first signs of insect infestation are observed, preferably before damage from feeding occurs. For best results, apply thoroughly to upper and lower leaf surfaces. Repeat application at 7 to 10 day intervals as needed.	
IN CALIFORNIA Lima Bean Pod Borer, Corn Earworm, Lygus and Stink Bug	10
Dust thoroughly upper and lower leaf surfaces.	
TOMATOES, EGG PLANTS, PEPPERS Tomato Fruitworm, Fall Armyworms, Flea Beetles, European Corn Borer, Stink Bugs	10
CUCUMBERS, SUMMER SQUASH Pickle and Melonworms, Striped and spotted Cucumber Beetles, Squash Bugs, Flea Beetles	10

NOTICE: This product is a restricted use pesticide. It may only be used by or under the direct supervision of a person licensed by the state in which it is used. For more information, contact your local agricultural agent or the manufacturer.

NET WT. 50 LBS.

THE CHAS. H. LILLY CO.
PORTLAND, OREGON

