



Miller's SEVIN* 5D

(5% SEVIN* DUST)

ACTIVE INGREDIENT

Carbaryl (1-naphthyl N-methylcarbamate)*..... 5%
 INERT INGREDIENTS 95%

*As the Reg. 744 of the Insecticide, Fungicide, and Rodenticide Act
 U.S. Pat. No. 2,903,474

E.P.A. Reg. No. 802-273-AA

CAUTION — Keep out of reach of children

HARMFUL IF INHALED OR SWALLOWED. Do not breathe dust. Do not take internally.

Avoid contact with skin. Wash hands before eating. Take shower or bath after work. Wear regular long sleeved work clothing. Change to clean clothing daily.

Note to Physician .. Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

Avoid contamination of foodstuffs, feed, feeding troughs, and watering receptacles. Keep out of reach of animals.

Dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes. Do not contaminate water by cleaning of equipment or disposal of wastes.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

PREHARVEST AND GRAZING USE LIMITATIONS

No post-treatment time limitations on Beans, Cucumbers, Eggplants, Peppers, Potatoes, Summer Squash, Sweet Corn, Tomatoes. When label directions for use are followed storage vines of Beans, Corn, Cucumbers and Summer Squash may be fed to meat and dairy animals.

Crop and Pest

DIRECTIONS

**Lbs
per acre**

CORN ... Corn Earworm 20 to 40

Start dusting when silks first appear. Repeat at 2 to 3 day intervals throughout silking period. Direct duster outlets toward ear zone to insure thorough coverage of each silk. (Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee population.)

POTATOES ... Colorado Potato Beetles, Leafhoppers, Flea Beetles 10 to 20*

Dust thoroughly upper and lower leaf surfaces. Do not graze treated fields or feed vines to dairy or meat animals.

BEANS ... Mexican Bean Beetles 10

Flea Beetles, Leafhoppers, Cucumber Beetles 20

Apply when first signs of insect infestation are seen, preferably before damage from feeding occurs. For best results apply thoroughly to upper and lower leaf surfaces. Repeat applications at 7 to 10 day intervals as needed.

IN CALIFORNIA ... Lima Bean Pod Borer, Corn Earworm, Lygus and Stink Bug 40

Dust thoroughly upper and lower leaf surfaces.

TOMATOES, EGGPLANTS, PEPPERS

Tomato Fruitworm, Fall Armyworms, Flea Beetles, European Corn Borer, Stink Bugs 20 to 40*

CUCUMBERS, SUMMER SQUASH

Pickle and Melonworms 10 to 20*

Striped and Spotted Cucumber Beetles, Squash Bugs, Flea Beetles 20*

*See application on appearance of insect. Repeat at 7 to 10 day intervals as needed.

NET WT 50 LBS.

THE CHAS. H. LILLY CO.
 PORTLAND, OREGON

