

The following brushy plants are controlled through a consistent year-to-year weed killing spray program:

Wild Blackberries	Poplar	Willows
Alder	Elderberry	Wild Rose
Birch	Raspberry	Polygon Ivy
Poison Oak	Salmonberry	

Apply Miller's BLACKBERRY and BRUSH KILLER to control brushy plants along fence rows, right-of-ways, and industrial areas where brushy plants are a problem.

SUMMER SPRAY—when brush first comes into full leaf
Dilution Chart and Application Directions

Use this amount of Miller's BLACKBERRY and BRUSH KILLER	Diluted with this amount of water
6 tablespoons	1 gallon
1 pint	6 gallons
1 quart	12 gallons

1. Spray when brush first comes into full leaf. (Summer Spraying—dilute with water only.)
2. Thoroughly wet entire brush, inside and outside, from tip to crown.
3. Soak crown thoroughly.
4. For best results, summer spray should be followed by Dormant Spray application as outlined below.


DORMANT SPRAY—after leaves have fallen in late fall or early winter
Dilution Chart and Application Directions

Use this amount of Miller's BLACKBERRY and BRUSH KILLER	Diluted with this amount of oil (Kerosene or Stove Oil)
1 pint	1 gallon
1 quart	2 gallons

1. Spray in late fall or winter. (For Dormant Spraying, dilute with oil only and use oil resistant hose.)
2. Thoroughly soak crowns and wet canes or trunks to a distance of 18 to 24 inches above the ground.
3. If growth is extremely heavy, it can be cut out and all crowns, stubs, and freshly cut surfaces sprayed to obtain complete coverage.
4. Spray all regrowth the following spring as outlined under Summer Spray.

NET CONTENTS
ONE QUART

\$4.69



CAUTION Keep out of the reach of children
See right side panel for additional cautions.

SPECIFIC

STUMP KILLING

1. Use 4 tablespoons (1 pint in 1 gal.).
2. Wet thoroughly all bark.
3. Stumps may be treated on freshly cut stumps.

KILLING STANDING TREES

1. Make a single girdle cut through the bark.
2. Liberally treat the cut and BRUSH KILLER, or other light oil (1 c).

Harmful if swallowed! Avoid spray drift to susceptible plants, fruit trees, etc. may injure them. Use coarse spray to drift. Use separate spray killing only. Do not contact domestic purposes. Vapors susceptible plants in the seeds, fertilizers, insecticides.

Do not use around the DO NOT USE, POUR, OPEN FLAME.

Do not graze dairy animals after application. Do not within 2 weeks of slaughter.

This product is toxic to ponds. Do not contaminate disposal of wastes.

Destroy empty containers.

NOTICE: Follow directions carefully. Keep containers tightly closed. Do not use in the vicinity of food. Do not use in the vicinity of food.

THE C
FOR
EPA

controlled through a consistent program:

Willows
Wild Rose
Poison Ivy

and BRUSH KILLER to control right-of-ways, and industrial problem.

NET CONTENTS
ONE QUART

\$469

not comes into full leaf
Application Directions

<p>Diluted with this amount of water</p> <p>1 gallon 6 gallons 12 gallons</p>

to full leaf. (Summer Spraying—
side and outside, from tip to crown

could be followed by Dormant Spray

have fallen in late fall or


Application Directions

<p>Diluted with this amount of oil (Diesel or Shovel Oil)</p> <p>1 gallon 2 gallons</p>

Dormant Spraying, dilute with oil

canes or trunks to a distance of

can be cut out and all crowns, tryed to obtain complete coverage, spring as outlined under Summer



CAUTION Keep out of the reach of children
See right side panel for additional cautions.

SPECIFIC DIRECTIONS

- STUMP KILLING** *SITE - UNSPECIFIED*
1. Use 4 tablespoons in one pint Diesel or other light oil (1 pint in 1 gal.).
 2. Wet thoroughly all exposed parts of stump and bark. Cut all regrowth around stump and treat newly cut stubs and bark.
 3. Stumps may be treated any time of year. Best results are had on freshly cut stumps two inches or more across.

- KILLING STANDING TREES** *SITE - UNSPECIFIED*
1. Make a single girdle of deep, overlapping axe or hatchet cuts through the bark at any convenient height.
 2. Liberally treat the injured area with Miller's BLACKBERRY and BRUSH KILLER, mixed 2 tablespoons per pint of Diesel or other light oil (1 cup in 1 gallon).

CAUTION

Harmful if swallowed! Avoid contact with skin, eyes or clothing. Avoid spray drift to susceptible plants such as flowers, vegetables, grapes, fruit trees, shrubs and ornamentals as this product may injure them. Use coarse droplet sprays as they are less likely to drift. Use separate sprayer or sprinkling can for weed or brush killing only. Do not contaminate water used for irrigation or domestic purposes. Vapors from this product may cause injury to susceptible plants in the immediate vicinity. Do not store near seeds, fertilizers, insecticides or fungicides.

Do not use around the home or recreation areas.
DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME.

Do not graze dairy animals on treated areas within 6 weeks after application. Do not graze meat animals on treated areas within 2 weeks of slaughter.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

Destroy empty container. Do not reuse for any purpose.

NOTICE: Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given hereon.

THE CHAS. H. LILLY CO.
PORTLAND, OREGON
EPA Est. 802-OR-1

48