


This is a facsimile of a pre-printed bag
on which the attached label is applied.



Net weight 50 lbs.

PERTHANE* 10D

(10% Perthane* Dust)

AGRICULTURAL INSECTICIDE

Active ingredients

Diethyl Diphenyl Dichloroethane	9.50%
Related Reaction Products	0.50%
Inert ingredients	90.00%

CAUTION— Keep out of reach of children

Avoid inhalation and contact with skin. In case of contact, wash thoroughly with soap and running water.

Avoid contamination of feed and foodstuffs. Do not use on feed or forage to be fed dairy animals or animals being finished for slaughter.

This product is toxic to fish. Keep out of lakes, streams, and ponds.

USE PRECAUTIONS

Do not apply closer to harvest than number of days indicated for specific crop being dusted.

Dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes.

*PERTHANE is the Reg. T. M. of the Rohm & Haas Co.

NOTICE: Follow directions carefully. Timing and method of application, weather and crop condition are factors with which user of chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risk of use, storage or handling of this material not in strict accordance with directions given herewith.

U.S.D.A. Reg. No. 802-291

MP-3F

GENERAL DIRECTIONS FOR FOLLOWING VEGETABLE CROPS ...

Apply dust when insects appear, making thorough applications. Repeat at weekly to 10 day intervals, or as required for adequate control.

SPECIFIC CROPS

CROP AND PEST	Lbs. Per Acre	Days Between Last Application and Harvest
<u>BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER, KOHLRABI</u> - Cabbage Worm, Cabbage Looper	25	3
<u>LETTUCE</u> - Leafhoppers, Cabbage Looper, Imported Cabbage Worm.	25	4
<u>SPINACH</u> - Leafhoppers, Cabbage Loopers	25	7
<u>CHERRIES</u> - Cherry Fruit Fly.	50	2

Make first application when adult flies appear, usually just after the blossoms open. A second application should be made after the shucks have fallen. Repeat every 10 days, or as required.

An application of Miller's PERTHANE 4E (spray) after harvest is suggested, especially if high temperatures have kept flies active.