


This is a facsimile of a pre-printed bag  
on which the attached label is applied.  
This label is the finished printed label  
as it will appear in this spot on the bag.



Net weight 50 lbs.

# G-THION 2D

Contains 2% GUTHION\*  
Store in a cool dry place  
AGRICULTURAL INSECTICIDE

Active ingredient, O,O-Dimethyl  
S-4-oxo-1,2,3-benzotriazin-3(4H)-  
ylmethyl phosphorodithioate\* ..... 2.0%  
Inert ingredients ..... 98.0%  
\*U.S. Pat. No. 2,758,115; Canadian Pat.  
No. 552,660. Reg. U.S. & Canadian Pat.  
Offs. by Farbenfabriken Bayer A.G.

## WARNING— Keep out of reach of children

May be fatal if swallowed, inhaled or absorbed through the skin. Do not get in eyes, on skin or on clothing. Do not breathe dust. Work to windward to stay out of dust. Wash thoroughly after handling. Wash all contaminated clothing with soap and hot water before reuse. Keep all unprotected persons out of the operating area or vicinity where there may be danger of drift.

Not for use or storage in or around the home. Vacated areas should not be reentered until the drifting insecticide and volatile residues have dissipated. Wear natural rubber gloves, protective clothing & goggles. Wear a mask or respirator of a type passed by the U. S. Bureau of Mines for Guthion protection.

Do not contaminate feed or foodstuffs. Keep out of reach of domestic animals.

TO PHYSICIAN: Atropine is antidotal. 2-TAM is also antidotal and may be administered in conjunction with Atropine.

## ENVIRONMENTAL CAUTIONS

This product is toxic to fish and wildlife. Keep out of lakes, streams and ponds. Birds feeding on treated areas may be killed. Do not apply to any area not specified on the label. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply when weather conditions favor drift of dust from areas treated. Do not apply where runoff is likely to occur.

## USE DIRECTIONS

Apply this product only as specified on this label. Do not use on other crops used

for food or forage. Application at rates above those shown may result in illegal crop residues. Do not treat food crops grown in the greenhouse with this product. Do not apply to crops in bloom or allow to drift to crops in bloom when pollinators are active.

Do not graze livestock in treated groves or orchards for 21 days after treatment.

Dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes.

## GENERAL DIRECTIONS

Apply the specified dosage for the specific crop as listed. Make first application when insects first appear and repeat as necessary. Observe all USE PRECAUTIONS and time limitations.

\* Do not apply closer to harvest than number of days indicated for specific crop being treated (shown in parentheses by each crop).

## FIELD AND VEGETABLE CROPS

ALFALFA, CLOVER (21\*) -- Use 30 to 35 lbs. per acre for Alfalfa Plant Bugs, Aphids, Leafhoppers, Lygus bugs, Mites.

BEANS, Snap (7\*) & Dry\* (30\*) -- Use 25 to 30 lbs. per acre for Aphids, Bean Leaf Beetles, Leafhoppers, Leaf Miners, Leaf Rollers and Mites. Use 30 lbs. per acre for Mexican Bean Beetles, Spotted or Striped Cucumber Beetles.

\* On Dry beans, do not apply more than 5 times per season. Do not feed or ensile treated forage.

BROCCOLI (1\*), BRUSSELS SPROUTS (7\*), CABBAGE (21\*), CALIFLOWER (15\*) -- Use 30 to 35 lbs. per acre for Aphids, Cabbage Loopers, Diamondback Moths, Imported Cabbage Worms.

POTATOES (7\*) -- Use 30 lbs. per acre for Colorado Potato Beetle. Use 30 to 35 lbs.

per acre for Aphids, European Corn Borer, Flea Beetles, Leafhoppers, Spittle bugs.

SPINACH (14\*) -- Use 25 to 35 lbs. per acre for Aphids, Leaf Miners, Mites.

TOMATOES (0\*) -- Use 25 lbs. per acre for Colorado Potato Beetle, Green Stink Bugs. Use 35 lbs. per acre for Tomato Fruitworm & Pinworm. Use 25 to 30 lbs. per acre for Drosophila, Leafminers. Use 30 to 35 lb. per acre for Aphids, Flea Beetles, Hornworms and Leafhoppers.

## FRUITS, BERRIES AND NUTS

BLACKBERRIES, BOYSENBERRIES, LOGANBERRIES, RASPBERRIES (14\*) --- Use 25 to 45 lbs. per acre for Leafhoppers, Leaf rollers, Obscure Root Weevil Adult.

GRAPES (10\*) -- Use 30 to 45 lbs. per acre for Grape Berry Moth, Leafhoppers, Mites. Use 35 to 45 lbs. per acre for Red-banded Leaf Roller. \*Do not apply more than 3 times per season.

STRAWBERRIES (5\*) --- Use 25 lbs. per acre for Meadow Spittlebug, Aphids, Omnivorous Leaf Tier (Strawberry Fruitworm).

WALNUTS (\*) --- Use 60 to 75 lbs. per acre for Aphids, Codling Moth, European Red Mite. \*Do not apply after husks split.

## ORNAMENTAL AND SHADE TREES IN THE NURSERY,

AND FOREST TREES -- Use 30 lbs. per acre for Aphids, Lace Bugs, Leafhoppers, Mites and Thrips. Use 25 to 35 lbs. per acre for European Pine shoot Moths and time applications to coincide with moth flight. Thorough coverage is necessary for best results.

Do not allow children or pets on treated area until the material has been washed into the soil and treated area is dry.

NOTICE: This product is registered under E.P.A. Reg. No. 802-310-AA MP-2F