

Miller's DIAZINON 4D

(4% Diazinon Dust)

ACTIVE INGREDIENT

*O,O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate 4.0%

INERT INGREDIENTS 96.0%

*Diazinon is the Reg. T.M. of CIBA-GEIGY
E.P.A. Reg. No. 802-318-AA

CAUTION - Keep out of reach of children

Harmful if swallowed. Avoid breathing dust. Avoid contact with skin, eyes or clothing. Wash thoroughly after handling.

Avoid contamination of feed and food products.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Destroy empty container by burying in non-crop lands away from water supplies or by burning (stay out of smoke).

ENVIRONMENTAL CAUTIONS

This product is toxic to birds, fish, and wildlife. Birds and other wildlife in treated areas may be killed. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

GENERAL DIRECTIONS

Apply the specified dosage for the specific crop as listed. Make first application when insects first appear and repeat as necessary.

NOTICE: Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

MP-4J

Crop and Pest

°Days Pounds per acre

BEANS (Snap, Pole and Lima) - Aphids, Spider Mites, Nitidulid Beetles, Western Spotted Cucumber Beetles.....	7	35
After harvest is complete, treated forage may be fed to livestock after 1 day following treatment, or cut for hay after 4 days following treatment.		
BROCCOLI, CABBAGE, CAULIFLOWER - Aphids, Diamondback Moth, Imported Cabbageworm	*	40
*5 days on Broccoli and Cauliflower; 7 days on Cabbage.		
CUCUMBERS - Aphids, Striped Cucumber and Western Spotted Cucumber Beetles.....	7	40
HOPS - Hop Aphid, Spider Mites, Western Spotted Cucumber Beetle, Beet Armyworm, Corn Earworm, Cutworms	14	50
KALE, COLLARDS - Aphids, Diamondback Moth	10	40
MELONS (Cantaloupe, Watermelon) - Aphids, Leafhoppers, Serpentine Leafminers	3	30-35
ONIONS (Green or Dry) - Pea Leafminer, Thrips	10	30
PARSLEY - Aphids	12	40
PEPPERS - Aphids, Serpentine Leafminer	5	40
SQUASH (Summer, Winter) - Aphids, Leafhoppers, Serpentine Leafminers, Thrips	*	35
*3 days on Winter; 7 days on Summer.		
STRAWBERRIES - Strawberry Aphids, Strawberry Leafrollers, Spittlebugs	5	30
SWISS CHARD - Aphids, Serpentine Leafminer, Western Spotted Cucumber Beetle	12	40
TOMATOES - Aphids, Banded Cucumber Beetles, Serpentine Leafminers	3	40

*Do not apply closer to harvest than number of days indicated for specific crop being treated.

NET WT. 50 LBS.

THE CHAS. H. LILLY CO.

PORTLAND, OREGON
EPA Est 02-OR-1

