

LEAD ARSENATE-SULPHUR 1585D

(15% Commercial Lead Arsenate - 85% Commercial Sulphur Dust)

Active ingredients

Lead Arsenate	14.1%
Sulphur	83.3%
Inert ingredients.	2.6%
Arsenic, expressed as metallic	3.045%
Arsenic in water soluble form, expressed as metallic.	0.075%
Lead, expressed as metallic . . .	3.400%

WARNING— Keep out of reach of children

May be Fatal if Swallowed! Avoid contact with skin, eyes or clothing. Wash thoroughly after using. Avoid breathing dust.

Dispose of wastes by burying in non-crop lands away from water supplies. The containers should be disposed of by punching holes in them and burying with wastes, or by burning (stay out of smoke).

USE PRECAUTIONS

Do not use sulphur during period of high temperatures. Do not graze or feed contaminated cover crops to livestock. Avoid drift to adjoining food and forage crops.

CHEERRY FRUIT FLY AND BROWN ROT

CHEERRIES FOR PROCESSING OR FRESH MARKET

1. Use 50 pounds per acre.
2. For Brown Rot Blossom Blight, make 3 applications during blossoming period. Increase to 4 or 5 applications during wet weather.
3. For Cherry Fruit Fly, apply at 10 day intervals. Make the first application when flies emerge.
4. Dust to obtain complete coverage of trees.
5. Do not apply within 2 days of harvest.
6. For Brown Rot on Fruit, when combination dust is not required, apply Sulphur alone at the rate of 50 lbs. per acre. Sulphur alone may be used up to harvest. For timing and number of sulphur applications, consult State Agricultural Experiment Station or State Extension Service Authorities.