

**MILLER'S
SLUG, SNAIL AND INSECT KILLER BAIT**

Can be used around vegetable gardens

KILLS: Slugs, Snails, Ants, Earwigs, Cutworms, Armyworms, Sowbugs, Millipedes, Crickets, Grasshoppers

PROTECTS: Flowers, Ornamentals, Shrubs and certain listed vegetables
Remains effective after rain or sprinkling

ACTIVE INGREDIENTS:

Metaldehyde (2,4,6,8-tetramethyl-1,3,5,7-tetraoxycyclo-octane	2.0%
Carbaryl (1-naphthyl N-methylcarbamate)	5.0%
INERT INGREDIENTS	93.0%

Keep out of reach of children

CAUTION

This pesticide may be fatal to dogs or other pets if eaten. Keep pets out of treated area. Read and follow all directions. See side panel for additional precautionary statements.

NET WEIGHT 2 LBS. (908 g)

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: May be harmful if swallowed. Avoid breathing dust. Avoid contact with skin, eyes, or clothing. Wash thoroughly after use. Keep out of reach of children and domestic animals. Avoid contamination of feed and foodstuffs.

IMPORTANT: Keep away from children, dogs and pets. Bait may be attractive to dogs. Confine pets during application to prevent them from believing they are being fed. Use and store under conditions where there will not be a hazard.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Drink one to two glasses of water and induce vomiting by touching back of throat with finger. Get medical attention. **IF INHALED:** Remove to fresh air. If person is not breathing give artificial respiration, preferably mouth-to-mouth. Get medical attention. **IF IN EYES:** Flush eyes with plenty of water. Call a physician if irritation persists. **IF ON SKIN:** Wash with plenty of soap and water.

ENVIRONMENTAL HAZARDS

Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters. This pesticide may be fatal to dogs or other pets if eaten. Keep pets out of treated area.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. For household use only.

STORAGE AND DISPOSAL

Store this product in its original container and keep in a locked storage area out of reach of children and domestic animals. Do not reuse container. Securely wrap container in several layers of newspaper and discard in trash.

HIGHLY EFFECTIVE MEAL BAIT FOR DESTRUCTION OF . . .

Slugs, Snails, Ants, Earwigs, Cutworms, Sowbugs, Millipedes, Crickets, Grasshoppers, Armyworms

Protects Flowers, Ornamentals, Shrubs and certain listed Vegetables.

Remains effective even after rain or sprinkling.


Draws pests to it for better results.

Works two ways - by simple contact or by being eaten.

SLUG, SNAIL AND INSECT KILLER BAIT is ready-to-use and remains effective even after rains and sprinklings. Protects flowers, ornamentals, shrubs and certain listed vegetables.

Be sure area to be baited is damp. In dry weather, water area well before baiting. Apply bait in evening or at night as these pests feed mainly at night. Until under control, bait may be reapplied as often as indicated for each type of plant.

Slugs and Snails favor dark, damp places. Bait especially well shaded areas around flowers, shrubs, rockeries, and other ornamentals where these pests stay during day-light hours.



Slugs, Snails and the listed insects are attracted to the bait. Place bait upwind to the infestation for best results. Upwind is the direction from which the wind is blowing from bait toward pests.

Slugs, Snails and the listed insects travel or migrate extensively. When invasions from adjacent areas are a probability, a lightly applied band of Slug, Snail and Insect Killer Bait the entire length of the infested areas will provide a barrier and aid in control.

1 pound of bait = 4 cups.

FLOWERS, ORNAMENTALS, SHRUBS

To kill slugs, snails, ants, earwigs, cutworms, sowbugs, millipedes, crickets and grasshoppers. Broadcast the bait lightly over the infested area using 1 lb. per 250 sq. ft. (6 Tblsp per 25 sq. ft.).

VEGETABLES AND FRUIT - To kill slugs, snails, armyworms, cutworms, earwigs, sowbugs, crickets, grasshoppers. Broadcast lightly on soil surface around plants at the rate indicated for each crop. Do not allow bait to contact foliage or edible part of crops. Observe the number of days before harvest for last application as noted in parentheses following each crop.

Asparagus (1): Use 4 tsp per 25 sq. ft. Reapply as often as every 7 days when applications are to spears; apply as often as every 3 days when applications are to ferns. Make no more than 3 applications prior to harvest and 2 additional applications to ferns only.

Snap Beans (fresh), Peas (fresh) (all 3): Use 1 Tblsp per 25 sq. ft. Reapply as often as every 7 days. Make no more than 4 applications per year.

Beets (roots), Carrots, Radish, Parsnips, Rutabaga, Turnips (roots) (7): Use 4 tsp per 25 sq. ft. Reapply as often as every 7 days. Make no more than 3 applications per year.

Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Kohlrabi (3): Use 4 tsp per 25 sq. ft. Reapply as often as every 7 days. Make no more than 3 applications per year.

Chinese Cabbage, Kale, Collards (14): Use 4 tsp per 25 sq. ft. Reapply as often as every 7 days. Make no more than 3 applications per year.

Sweet Corn (2): Use 4 tsp per 25 sq. ft. Reapply as often as every 3 days. Make no more than 8 applications per year.

