

GENERAL DIRECTIONS

SOIL APPLICATION

prior to planting the following vegetables:

Beans	Cucumbers	Peas
Broccoli	Endive	Peppers
Cabbage	Lettuce	Potatoes
Carrots	Melons	Spinach
Cauliflower	Onions	Squash
Corn	Parsnips	Tomatoes

To control Cutworms, Seed-corn Maggots, Wireworms

Apply evenly as a broadcast treatment over the surface of the soil. The amount in this package (1 lb) should be broadcast over 450 square feet. Immediately after spreading, mix thoroughly with the soil to a depth of 6 to 8 inches.

SPECIAL USES

(based on 12-inch row spacing)

Onions (onion maggot): apply in furrow at the rate of 6 tablespoons per 100 feet of row at time of planting.

Radishes (root maggot): apply in furrow at rate of 9 tablespoons per 100 feet of row at time of planting.

CAUTION

Harmful if swallowed. May be absorbed through skin.

Avoid breathing dust. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling.

Avoid contamination of feed and foodstuffs.

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Destroy empty container—do not reuse for any purpose.

NOTICE: Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material, not in strict accordance with directions given hereon.

THE CHAS. H. LILLY CO.
PORTLAND, OREGON
EPA Est. 802-OR-1

NET WT.
1 LB.



MULTI-USE Diazinon[®] Insect dust

FOLIAGE INSECTICIDE FOR VEGETABLES • FRUITS • FLOWERS

ACTIVE INGREDIENT

0,0-diethyl 0-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate 4.0%

INERT INGREDIENTS 96.0%
Total 100%

*DIAZINON® is the Reg. T.M. of
CIBA-GEIGY

EPA Reg. No. 802-360-AA

CAUTION

Keep out of the reach of children
See left side panel for additional cautions.

Miller's
MULTI-USE

Diazinon Insect Dust GENERAL DIRECTIONS

FOLIAGE APPLICATIONS

Vegetables, Flowers, Fruits

- 1 Dust lightly on first appearance of insect or insect damage
- 2 Make complete coverage of all pests and foliage
- 3 Repeat applications as necessary for control, but observe the number of days before harvest for last application noted by each crop (for instance, on Beans, you may apply this dust up to 7 days before harvest or picking)

BEANS (7), **PEAS** (day of harvest), **POTATOES** (35), **TOMATOES** (3)—Aphids, Leafhoppers.

BEETS (14), **BROCCOLI** (5), **CABBAGE** (7), **CAULIFLOWER** (5), **CARROTS** (10), **RADISHES** (10)—Aphids.

ONIONS (10), **CUCUMBERS** (7), **SUMMER SQUASH** (7), **WINTER SQUASH** (3), **MELONS** (3)—Aphids, Thrips, Leaf Miners, Mites.

APPLES (14), **PEARS** (14)—Aphids, Mites.

STRAWBERRIES (5)—Aphids, Cyclamen and Two-Spotted Mites.

ROSES, FLOWERS, SHRUBS, ORNAMENTALS—Aphids, Mites, Leafhoppers, Mealy Bugs, White Fly, Scale Crawlers, Lace Bug.

USE PRECAUTIONS

Do not use on certain ferns, poinsettias, hibiscus and gardenias since injury may occur. Not effective against certain spider mites on evergreens.

Do not apply when honey bees are foraging in the garden.

GENERAL DIRECTIONS

NET WT.
1 LB

SOIL APPLICATION

Beans	Chickpeas	Leas
Broccoli	Endive	Peppers
Cabbage	Lettuce	Potatoes
Carrots	Melons	Spinach
Cauliflower	Onions	Squash
Corn	Parsnips	Tomatoes

To control Cutworms, Seed-corn Maggots, Wireworms

Apply evenly as a broadcast treatment over the surface of the soil. The amount in this package (1 lb.) should be broadcast over 450 square feet. Immediately after spreading, mix thoroughly with the soil to a depth of 6 to 8 inches.

SPECIAL USES

(based on 12-inch row spacing)

Onions (onion maggot): apply in furrow at the rate of 6 tablespoons per 100 feet of row at time of planting.

Radishes (root maggot): apply in furrow at rate of 9 tablespoons per 100 feet of row at time of planting.

CAUTION

Harmful if swallowed. May be absorbed through skin.

Avoid breathing dust. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling.

Avoid contamination of feed and foodstuffs.

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Destroy empty container - do not reuse for any purpose.

NOTICE: Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material, not in strict accordance with directions given herewith.

THE CHAS. H. LILLY CO.
PORTLAND, OREGON
EPA Est. 802-OR 1



MULTI-USE Di Insect

FOLIAGE INSECT
VEGETABLES • FRUITS

ACTIVE INGREDIENT
0,0-diethyl 0-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate
INERT INGREDIENTS
Total

ACCEPTED
11-21-74
EPA Reg. No. 100-100-01
CIBA-GEIGY

CAUTION Keep out of reach of children. See left side panel for first aid instructions.