

*Miller's*

# TRIBASIC COPPER SULPHATE 3D

A FUNGICIDAL DUST

FOR CONTROL OF FIRE BLIGHT OF PEARS

DURING BLOSSOMING PERIOD

E.P.A. REG. No. 802-364-AA

**ACCEPTED**

**JUN 27 1973**

UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENT CIDE ACT  
FOR ECONOMIC POISON REG STER-  
ED UNDER NO. 802-364 SUBJECT  
TO ATTACHED COMMENTS.

**Active ingredient:**

Copper (from Tribasic Copper Sulphate),  
expressed as metallic ..... 3.0%  
Inert ingredients ..... 97.0%

**CAUTION — Keep out of reach of children.**

Avoid breathing dust. Avoid contact with skin,  
eyes or clothing. Wash thoroughly after using.

Avoid contamination of feed and foodstuffs.

Dispose of wastes by burying in non-crop lands  
away from water supplies. Containers should be  
disposed of by burying with wastes.

This product is toxic to fish. Keep out of lakes,  
streams or ponds. Do not apply where runoff is  
likely to occur. Do not apply when weather con-  
ditions favor drift from areas treated. Do not  
contaminate water by cleaning of equipment, or  
disposal of wastes. Apply this product only as  
specified on this label.

**DIRECTIONS FOR USE**

PEARS - Fire Blight Control

1. Apply 25 pounds of dust per acre.
2. Make first application within three days  
after first blossoms open (30% full bloom).
3. Repeat at 5 day intervals through blossoming  
period. (Keep blossoms covered to protect  
through period of high humidity.)
4. Do not apply copper to Anjous or Seckels. or  
to trees when they are wet.

**NOTICE:** Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

NET WT. 50 LBS.

THE CHAS. H. LILLY CO.  
PORTLAND, OREGON

MP=4H