

# Miller's MCP SODIUM SALT 2

## GENERAL DIRECTIONS

Mix with water only. Add one-half required amount of water to spray tank, then add specified amount of this product with agitation, finally adding the balance of the water with continued agitation. Apply during warm weather when weeds are growing actively.

## SPECIFIC DIRECTIONS

**RICE**—for the control of Water Plantain, Arrowhead Lily, Red Stem, and certain susceptible sedges.

1. Use 3 to 5 pints in 10 to 30 gallons of water per acre.
2. Apply 55 to 65 days after planting when Rice is full tillered and 6 to 8 inches above water, or when crop stems begin to elongate. Do not apply after Rice is in boot stage.

**FLAX**—For control of Mustard, Lambsquarters, and other susceptible weeds

1. Use 2 pints per acre in 20 to 70 gallons of water by ground equipment; or 10 gallons of water by air.
2. Apply when crop is 4 to 8 inches tall but before bud stage.

## WHEAT, OATS AND BARLEY

For control of Mustard, Lambsquarters, and other susceptible weeds.

1. Use 2 to 4 pints per acre in 5 to 20 gallons of water by ground equipment, or 2 to 3 gallons water by air. For Oats, use 2 pints per acre.
2. Apply from full tiller to boot stage. Do not spray from boot to dough stage.

NOTICE: Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

MP-4H

# Miller's MCP SODIUM SALT 2

**PEAS**—For control of Mustard, Lambsquarters and other susceptible weeds.

1. Use ½ to 1 pint in 20 to 60 gallons of water per acre. (For aerial application use 10 gallons total spray solution per acre.) Adjust dosage rates to air temperatures as follows:  
Less than 55°F—1 pint per acre  
55°F to 70°F—4/5 pint per acre  
70°F to 80°F—½ pint per acre
2. Apply after weeds emerge and before they are 3 inches tall. Do not apply later than the 3 node stage of growth of the peas before blooming period.
3. Do not apply to peas that are stressed from lack of moisture.
4. Do not feed treated vines to livestock. Do not graze or forage treated pea fields.

**NOTE:** Consult State Extension Service personnel or cannery company fieldmen for proper timing of applications with regard to varietal differences of growth rate.

Insist on...

# Miller's

# MCI SODIUM S

(contains 2 lbs. MCP acid equiv)

For selective control of Broadl  
PEAS - FLAX - RICE - OAT  
small Cereals

Active ingredient  
\*Sodium Salt of 2-Methyl-4-Chlorophenoxy  
Inert ingredients .....  
\*Equivalent to 21% 2-Methyl-4-Chlorophenoxy

E.P.A. Reg. No. 802-374

USE ONLY IN ACCORDANCE WITH  
RECOMMENDATIONS OR RES

## WEED KILLER

**CAUTION** — Keep out of reach

See right side panel for additional c

NET CONTENTS: \_\_\_\_\_

THE CHAS. H. LILLY CO.  
PORTLAND, OREGON

*Miller's*  
**MCP**  
**SODIUM SALT 2**

**PEAS**—For control of Mustard, Lambsquarters and other susceptible weeds.

1. Use ½ to 1 pint in 20 to 60 gallons of water per acre. (For aerial application use 10 gallons total spray solution per acre.) Adjust dosage rates to air temperatures as follows:  
 Less than 55°F—1 pint per acre  
 55°F to 70°F—4/5 pint per acre  
 70°F to 80°F—½ pint per acre
2. Apply after weeds emerge and before they are 3 inches tall. Do not apply later than the 3 node stage of growth of the peas before blooming period.
3. Do not apply to peas that are stressed from lack of moisture.
4. Do not feed treated vines to livestock. Do not graze or forage treated pea fields.

**NOTE:** Consult State Extension Service personnel or cannery company fieldmen for proper timing of applications with regard to varietal differences of growth rate.



**MCP**  
**SODIUM SALT 2**

(contains 2 lbs. MCP acid equivalent per gallon)

**For selective control of Broadleaved Weeds in  
 PEAS - FLAX - RICE - OATS and other  
 small Cereals**

Active ingredient  
 \*Sodium Salt of 2-Methyl-4-Chlorophenoxyacetic acid...23.2%  
 Inert ingredients .....76.8%  
 \*Equivalent to 21% 2-Methyl-4-Chlorophenoxyacetic acid

E.P.A. Reg. No. 802-374-AA

**USE ONLY IN ACCORDANCE WITH LOCAL AND STATE  
 RECOMMENDATIONS OR RESTRICTIONS**


**WEED KILLER**

**CAUTION — Keep out of reach of children**

See right side panel for additional cautions

**NET CONTENTS:**

**GALLONS**



*Miller's*  
**MCP**  
**SODIUM SALT 2**

**CAUTION**

**PLEASE READ ENTIRE LABEL. USE ONLY AS DIRECTED, NOTING ALL CAUTIONS AND WARNINGS.**

Harmful if swallowed.

This product may cause skin irritation. Avoid contact with skin, eyes and clothing. In case of contact, wash skin with plenty of soap and water; flush eyes with plenty of water for at least 15 minutes and get medical attention. Keep out of reach of children and domestic animals.

Do not store with foodstuffs. Do not transport or store near fertilizers, insecticides, fungicides or seeds.

Do not apply directly to or otherwise permit sprays to come in contact with vegetables, flowers, grapes, fruit trees, cane berries, ornamentals or other desirable plants which are sensitive to MCP and do not permit spray mist to drift onto them, since even minute quantities of the spray may cause severe injury during both growing and dormant periods. Coarse sprays are less likely to drift.

Application by airplane, ground equipment, or hand sprayers should be made only when there is no hazard from drift.

Do not apply by airplane in the vicinity of grapes or other desirable MCP susceptible plants.

Do not contaminate irrigation ditches or water used for domestic purposes. Excessive amounts of MCP in the soil may temporarily inhibit seed germination or plant growth.

At higher temperatures vaporization may cause injury to susceptible plants growing nearby. Keep container tightly closed when not in use.

Do not use spray equipment for any other purpose. Such equipment should not be used for application of insecticides, fungicides, or other agricultural chemicals.

Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes.

THE CHAS. H. LILLY CO.  
 PORTLAND, OREGON

*Miller's*