

Miller's
MCP
SODIUM SALT 2
(2 lbs. MCP acid equivalent per gallon)

ACTIVE INGREDIENT

*Sodium Salt of 2-Methyl-4-Chloro-
phenoxyacetic acid 23.2%

INERT INGREDIENTS 76.8%

*Equivalent to 21% 2-Methyl-4-
Chlorophenoxyacetic acid

WEED KILLER

For selective control of Broadleaved Weeds in
PEAS, FLAX, RICE, OATS, WHEAT, BARLEY,
ESTABLISHED PASTURES, TURFS AND OTHER
NON-CROP AREAS.

ACCEPTED
with comments
NOV 25 1985
Federal Insecticide, Fungicide, and Rodenticide Act
EPA Reg. No. 802-374

KEEP OUT OF REACH OF CHILDREN

DANGER

STATEMENT OF PRACTICAL TREATMENT

In case of contact, wash skin with plenty of soap and water; for eyes, flush with water for 15 minutes. Get medical attention. If swallowed call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat. Do not induce vomiting or give anything by mouth to an unconscious person.

See left side panel for Additional Precautionary Statements.

NET CONTENTS: GALLON(S)
The Chas. H. Lilly Co., Portland, Oregon 97218
EPA Est. 802-OR-1 EPA Reg. No. 802-374

Label size: 21 3/16 x 9 1/8 inches
Label Color: Black print on a white and
yellow front panel and a yellow-orange
striped side and back panel.

PRECAUTIONARY STATEMENTS

**HAZARD TO HUMANS AND
DOMESTIC ANIMALS**

DANGER

Corrosive. Causes eye and skin irritation. Do not get in eyes, on skin or clothing. Wear goggles or face shield and rubber gloves when handling. Harmful or fatal if swallowed. Do not breathe spray mist. Wash thoroughly after handling.

ENVIRONMENTAL HAZARDS

Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply when weather conditions favor drift from target areas. Avoid use of small-diameter spray nozzles.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

STORAGE: Do not store near other pesticides or seeds. Open dumping is prohibited.

PESTICIDE DISPOSAL: Pesticide, spray mixture or rinsate that cannot be used according to label instructions must be disposed of according to Federal, State or local procedures under the Resource Conservation and Recovery Act.

CONTAINER DISPOSAL: Triple rinse or equivalent and offer for recycling or reconditioning, or dispose of in a sanitary landfill, or by other approved state or local procedures.

NOTICE: Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

LC-2R

BEST AVAILABLE COPY

MIXING DIRECTIONS

Mix with water only. Add one-half required amount of water to spray tank, then add specified amount of this product with agitation, finally adding the balance of the water with continued agitation. Apply during warm weather when weeds are growing actively.

APPLICATION DIRECTIONS

Do not apply directly to or otherwise permit sprays to come in contact with vegetables, flowers, grapes, fruit trees, caneberries, ornamentals or other desirable plants which are sensitive to MCP and do not permit spray mist to drift on to them, since even minute quantities of the spray may cause severe injury during both growing and dormant periods. Coarse sprays are less likely to drift.

Do not apply by airplane in the vicinity of grapes or other desirable MCP susceptible plants.

Do not contaminate irrigation ditches or water used for domestic purposes. Excessive amounts of MCP in the soil may temporarily inhibit seed germination or plant growth.

At higher temperatures vaporization may cause injury to susceptible plants growing nearby. Keep container tightly closed when not in use.

Do not use spray equipment for any other purpose. Such equipment should not be used for application of insecticides, fungicides, or other agricultural chemicals.

SPECIFIC DIRECTIONS

RICE - for the control of Water Plantain, Arrowhead Lily, Red Stem and certain susceptible sedges.

1. Use 3 to 5 pints in 10 to 30 gallons of water per acre.
2. Apply 55 to 65 days after planting when Rice is full tillered and 6 to 8 inches above water, or when crop stems begin to elongate. Do not apply after Rice is in boot stage. Do not spray when temperature is over 90° F.

NOTE: Do not use in rice fields where shellfish are of economic importance. Do not grow crayfish or catfish in treated rice fields.

FLAX - for control of Mustard, Lambsquarters, and other susceptible weeds.

1. Use 2 to 4 pints per acre in 20 to 70 gallons of water by ground equipment; or 10 gallons of water by air.
2. Apply when crop is 4 to 8 inches tall but before bud stage.

WHEAT, OATS and BARLEY - for control of Mustard, Lambsquarters, and other susceptible weeds.

1. Use 2 to 4 pints per acre in 5 to 20 gallons of water by ground equipment, or 3 or more gallons water by air. For OATS, use 2 pints per acre. Apply from full tiller to boot stage. Do not spray from boot to dough stages.
2. For less susceptible weeds, use up to 6 pints per acre after crop has tillered and up to early boot stage. The 6 pint per acre application rate can produce injury to wheat.

BEST AVAILABLE COPY

GRASSES GROWN FOR SEED

1. Use 2 to 4 pints per acre in sufficient water to give adequate coverage. Use higher rate where weed stands are heavy.
2. In established grasses, apply in early spring or in the fall prior to the time seed heads appear in the boot. On new seedling grass, apply after grass has 5 leaves.

NOTE: MCP may injure bentgrass at the higher rates. Do not graze dairy animals in treated areas until 7 days after treatment. Do not graze meat animals in treated areas within 7 days of slaughter.

ESTABLISHED GRASSLANDS AND PASTURES

1. Use 2 to 6 pints per acre in sufficient water (10 to 100 gallons) to give thorough coverage. Use higher rate for Canada Thistle and other hard-to-kill weeds.
2. Spray perennials in bud stage and regrowth in fall; other weeds in spring or fall.

NOTE: Alfalfa, Alsike, Red, Crimson, Sub-clovers are injured by MCP.

ESTABLISHED LAWN AND TURFS

1. Use two fluid ounces in 3 to 5 gallons of water, apply uniformly over 1,000 sq. ft. For larger areas, use 2 to 6 pints per acre in 10 to 100 gallons of water.
2. Apply spring or fall when weeds are actively growing. Do not mow within two days before or after application.

NON-CROP SPRAYING - Canada Thistle, White Top, Meadow Buttercup, Field Bindweed (Morning-glory).

1. For spot treatment, use ½ pint in 3 to 4 gallons of water or 1½ gallons per acre in sufficient water to give coverage. Spray to wet weeds thoroughly.
2. Apply when weeds are in bud to early bloom and again on fall regrowth.

NOTE: Do not forage or graze meat or dairy animals on treated areas within 7 days of treatment.

BEST AVAILABLE COPY

5 9 5
NOTE: Do not forage or graze meat animals on treated areas within 7 days of slaughter. Do not forage or graze dairy animals on treated areas within 7 days after treatment. ✓

PEAS - For control of Mustard, Lambsquarters and other susceptible weeds.

1. Use $\frac{1}{2}$ to 1 pint in 20 to 60 gallons of water per acre. For aerial application use 10 gallons total spray solution per acre. Adjust dosage rates to air temperatures as follows: ✓

Less than 55° F — 1 pint per acre
55° F to 70° F — $\frac{4}{5}$ pint per acre
70° F to 80° F — $\frac{1}{2}$ pint per acre

2. Apply after weeds emerge and before they are 3 inches tall. Do not apply later than the 3 node stage of growth of the peas before blooming period. ✓
3. Do not apply to peas that are stressed from lack of moisture. ✓
4. Do not feed treated vines to livestock. Do not graze or forage treated pea fields. ✓

NOTE: Consult State Extension Service personnel or cannery company fieldmen for proper timing of applications with regard to varietal differences of growth rate. ✓

BEST AVAILABLE COPY