Label size: 47/8 x 19½ inches Front panel size: 47/8 x 57/8 inches

Front Panel:

MILLER'S ALPHASPRA 800

e.

1

· · · ·

· · ·

*1-Naphthaleneacetic Acid equivalent 19.40% or 800 grams per gallon.

A Hormone-like Spray Concentrate containing 800 grams 1-Naphthaleneacetic acid equivalent per gallon

USED TO REDUCE PREHARVEST DROP OF APPLES AND PEARS AND TO DELAY LEAF DROP OF CUT HOLLY

KEEP OUT OF REACH OF CHILDREN		>	10++
CAUTION -			14 6-1
See back panel for additional precautionary statemen	ts.		

NET CONTENTS ONE GALLON

THE CHAS. H. LILLY CO. PORTLAND, OREGON 97218 EPA Est. 802-OR-1 EPA Reg. No. 802-377

ACCEPTED

NOV 1 9 1982 Under the Federal Insection Fungatile, and Redenticide as amended, for the pesticide registered under, 2-377

Left Panel

PRECAUTIONARY STATEMENTS Hazards to Humans & Domestic Animals CAUTION

Harmful if swallowed. Avoid breathing spray mist. Avoid contact with skin, eyes and clothing. In case of contact, immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists.

ENVIRONMENTAL HAZARD

Keep out of lakes, ponds or streams. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

STORAGE: Avoid contamination of feed and foodstuffs. DISPOSAL: Triple rinse container and offer for recycling or reconditioning, or dispose of in a sanitary landfill, or by incineration if permitted by state and local authorities.

NOTICE: Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assume all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

LC-6Q

Right Panel-

Miller's ALPHASPRA 800

Miller's ALPHASPRA 800 is used to reduce pre-harvest drop of certain varieties of Apples and Pears and to delay leaf drop on cut Holly.

APPLES AND PEARS - Application Directions

Consult State Agricultural Experiment Station or State Extension Service Authorities for correct timing of application on Apples and Pears.

Spray trees thoroughly, wetting all fruit stems and adjacent leaves. Apply to Apples 7 to 14 days before harvest and to Pears 6 to 7 days before harvest. Do not spray within 5 days of harvest. Do not make more than 2 applications and do not delay picking beyond optimum maturity.

APPLES AND PEARS – Dilution for spraying with ground equipment Apples--Mix at the rate of 1.2 fl. oz. per 100 gallons of water (3 fl. oz. per 250 gals.) to prepare a 20 ppm spray solution. Pears-Mix at the rate of 0.6 fl. oz. per 100 gallons of water (1½ fl. oz. per 250 gals.) to prepare a 10 ppm spray solution.

Thoroughly cover trees with appropriate spray, using up to 465 gallons per acre on Apples of the 20 ppm spray solution which provides a naphthaleneacetic acid equivalent to 35 grams per acre. Use up to 670 gallons of 10 ppm spray per acre for Pears which provides 25 grams of naphthaleneacetic acid equivalent per acre.

HOLLY – to delay leaf drop

Use ¼ to ½ cupful to 100 gallons of water (equals 33 to 66 ppm). Dip Holly soon after cutting; do not soak; drain off excess solution. Pack before Holly dries with sufficient moisture to keep Holly fresh until it reaches its destination. Ship immediately or store at 34 to 40° F. Keep solution as clean as possible. Renew dip with fresh solution as it is used up. Do not keep solution longer than 3 to 4 days.