ALPHASPRA 800

ACTIVE INGREDIENT *1 Naphtheleneacetic acid, Potassium Salt INERT INGREDIENTS 100 00% fi-Naphthaleneacetic ecid equivalent 19 40% or 800 grams پنت gallon

A Hormone-like Spray Concentrate containing 800 grams 1-Naphthaleneacetic acid equivalent per gallon

USED TO REDUCE PREHARVEST DROP OF APPLES AND PEARS AND TO DELAY LEAF DROP OF CUT HOLLY

> Keep out of reach of children CAUTION

See back panel for additional precautionary statuments

ACCEPTED

DEC - 4 1984

Under the Federal Insection of Fungicide, and Rodentinide as amonded, for the pesticide registered under ERA Bag. No. 802

Lilly/Miller ALPHASPRA 800

Lilly Miller ALPHASPRA 800 is used to reduce preharvest drop of certain varieties of Apples and Pears and to delay leaf drop on cut Holly.

APPLES AND PEARS — Application Directories

Consult State Agricultural Experiment Station or State Extension Service Authorities for correct timing of application on Apples and

Spray trees thoroughly, wetting all fruit stems and adjacent leaves. Apply to Apples 7 to 14 days before harvest and to Pears 6 to 7 days before harvest. Do not spray within 5 days of harvest. Do not make more than 2 applications and do not delay picking beyond optimum maturity.

APPLES AND PEARS - Dilution for spraying with ground

Apples — Mix at the rate of 1.2 ft, oz. per 100 gallons of water (3 ft. oz. per 250 gals.) to prepare a 20 ppm spray solution. Pears -- Mix at the rate of 0.6 fl. oz. per 100 gallons of water (11/2 fl. oz.

per 250 gals.) to prepare a 10 ppm spray solution.

Thoroughly cover trees with appropriate spray, using up to 485 gallons per acre on Apples of the 20 ppm spray solution which provides a naphthaleneacetic acid equivalent of 35 grams per acre. Use up to 670 gallons of 10 ppm spray per acre for Pears which provides 25 grams of naphthaleneacetic acid equivalent per acre.

11

Prevention of Pre-Harvest Fruit Drop By Aerial Application

2

Alphaspra 800 becomes effective within 3-4 days of application and prevents drop for three or four weeks. Separate sprays should be applied to early and late maturing varieties.

Apples, D'Anjou Pears: Use 1/4 to ½ pt. in at least 5 gals water per acre. Apply 6-8 days before harvest. Do not make more than two applications per season. Do not apply within two days of harvest.

Pears, Bartlett, Bosc: Use 1/8 to ½ pt. in at least 5 gals per acre. Apply before drop begins, 6 to 8 days before harvest. Do not make more than two applications per season. Do not apply within two days of harvest. When Bartlett and Bosc are interplanted, spray only once. Use a maximum of ½ pt. per acre on Bartlett to prevent pre-mature ripening.

Aerial applications of Alphaspra 800 should be made under weather conditions favoring minimal drift to achieve a uniform distribution of product.

HOLLY -- To delay leaf drop

Use ¼ to ½ cupful to 100 gallons of water (equals 33 to 66 ppm). Dip Holly soon after cutting; do not soak; drain off excess solution. Pack before Holly dries with sufficient moisture to keep Holly fresh until it reaches its destination. Ship immediately or store at 34 to 40° F. Keep solution as clean as possible. Renew dip with fresh solution as it is used up. Do not keep solution longer than 3 to 4 days.

PRECAUTIONARY STATEMENTS Hazards to Humans & Domestic Animals CAUTION


Harmful if swallowed. Avoid breathing spray mist. Avoid contact with skin, eyes and clothing. In case of contact, immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists.

ENVIRONMENTAL HAZARD

Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.



STORAGE AND DISPOSAL

Do not contaminate water food, or feed by storage or disposal.

STORAGE: Store in a cool, dry, dark place.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.


NOTICE: Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals, not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

EPA Reg No 802-377

I C-ER

∳vailable size: 1 gal.

THE CHAS H LILLY CO.
PORTLAND, OREGON 97218
EPA Est. 802 OR 1



30/3