

Insist on...

Miller's



HEPTACHLOR 50W

(8 oz. actual Heptachlor per lb.)

**A Wettable Powder for use
in water as an Insecticidal Spray
and as a Seed Treater**

Active ingredients:	
Heptachlor*	50.0%
Related Compounds	19.4%
Inert ingredients:	30.6%
Total	100.0%

*Heptachlorotetrahydro-4,7-methanoindene

E.P.A. Reg. No. 802-379-AA

WARNING Keep out of the reach of children
See back panel for additional warnings.

May be fatal if swallowed! Do not breathe vapor, dust or spray mist. Do not get on skin. In case of contact, wash immediately with soap and water. Do not contaminate feed and foodstuffs.

This product is toxic to fish and wildlife. Keep out of lakes, streams and ponds. Birds feeding on treated areas may be killed. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Dispose of excess treated seeds by burial in non-crop lands away from water supplies.

Rinse equipment and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes or by burning (keep out of smoke).

NET WEIGHT 4 LBS.

THE CHAS. H. LILLY CO.
PORTLAND, OREGON

LILLY'S

Miller's

ACCEPTED

FEB 13 1974

ORDERED FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POLYMER REGISTER
AND UNDER NO. 802-379-AA

802-379

Miller's HEPTACHLOR 50W

Please read entire label.

Use only as directed, noting all Warnings and Use Precautions.

DIRECTIONS FOR USE

INSECTICIDAL SPRAY

Wash recommended amount of Miller's HEPTACHLOR 50W through screen into partially filled spray tank, or pre-mix in bucket half filled with water and add mixture into nearly filled spray tank. Add balance of water to obtain proper gallonage of completed spray solution. Do not allow mixture to stand. Keep agitator running during filling and spraying operations. Do not combine wettable powders with emulsifiable liquids in the same spray tank unless previous use of the materials being combined has proved to be compatible.

SOIL APPLICATIONS (Pre-plant treatment only)

Pre-plant soil applications may be used for the following crops. Mix the indicated amount of Miller's HEPTACHLOR 50W in sufficient water to assure uniform coverage over soil surface just before last discing when preparing seed bed. Follow application by tilling thoroughly to mix the chemical into the upper 3 to 6 inches of soil.

Crop	Insects	Lbs. per Acre
Corn	Wireworms	6 to 10
Corn	Corn Rootworm, Cutworms, Seed Corn Maggot, White Grub	4 to 6
Peppers	Wireworms	4 to 6
Tomatoes	Wireworms	4 to 6

PINEAPPLE—Ants

Use 1 to 2 lbs. per acre in sufficient water (200 to 300 gallons) per acre, to insure uniform coverage. Do not make application within 60 days of harvest.

FOLIAGE APPLICATIONS

LAWNS AND ORNAMENTAL TURFS

Ants, Asiatic Garden Beetle, White Grubs—Use 4 lbs. per acre of lawn or turf.

Chiggers, Fleas, Ticks—Use 2 lbs. per acre of lawn or turf.

Apply recommended dosage in sufficient water (50 or more gallons per acre) to insure uniform coverage. Be sure that the chemical has washed from grass into soil area and that grass is dry before allowing children or pets on lawn or turf. Do not feed clippings to livestock, or allow livestock to graze on treated turf.

SEED TREATER

Miller's Heptachlor 50W seed treatments are for the protection of germinating seeds and seedlings against the insects specified. Heptachlor 50W may be applied as a dust or slurry, and may be combined with commonly used fungicidal seed treatments. Treated seed may be bagged, marked: TREATED SEED. DO NOT USE FOR FOOD, FEED OR OIL PURPOSES, and stored until planting time. Slurry treated seed should not be stacked until it has had a chance to dry.

Mix the required amount with the seed in grain drill or planter, or add to the seed and thoroughly mix in a revolving or drum type treater prior to planting.

Rates of Application:	Ounces Per Bushel of Seed	Ounces Per 100 lbs. of Seed
WHEAT—Wireworms	1 1/4 to 1 3/4	2 to 3
OATS—Wireworms	2/3 to 1	2 to 3
CORN—Seed Corn Maggot	3/4 to 1	1 to 1 1/2

USE PRECAUTIONS—Seed Treater—Never permit treated seed to be used as food by humans or domestic animals.

Treated seeds exposed on soil surface will be hazardous to birds and other wildlife.

Dispose of excess treated seeds by burial in non-crop land away from water supplies.

NOTICE: Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

MP-4H