

DIRECTIONS FOR USE

ALFALFA

For control of Alfalfa Weevil, Alfalfa Caterpillar, Spittlebugs, Leafhoppers, Pea Leaf Weevil, Western Spotted Cucumber Beetle and Aphids. Use amount required in 10 gal. of water by air, or 20 gal. of water with ground equipment.

First Cutting — Apply 2 to 3 quarts per acre as soon as weevil damage appears in the field. Early application is desired to control the weevil when they are small. Do not apply within 7 days of harvest. Do not apply after full bloom.

Stubble Spraying — in areas where weevil appears at cutting time. Remove first cutting and apply 2 to 3 qts. per acre to stubble to control weevil larvae and aphids.

Second Cutting — Apply 2 to 3 qts. per acre to control alfalfa weevil and other insects as they appear. Do not apply within 7 days of harvest. Do not apply after full bloom.

CLOVER

For the control of **Clover Leaf Weevil**, apply 2 qts. (in approximately 25 gal. of water) per acre. Apply in spray when clover is 2 to 6 inches tall.

CLOVER—Continued

For the control of **Meadow Spittlebug**, apply 2 qts. (in approximately 25 gal. of water) per acre. Apply in spring within a week after eggs begin to hatch. As a fall treatment, apply 3 qts. (in approximately 25 gal. of water) per acre. Make application in early September.

Treat clover in bloom only when bees and other pollinators are not visiting plants. Do not apply closer than 7 days of harvest.

PEAS

For the control of **Pea Aphid**, **Pea Leaf Weevil**, **Alfalfa Looper**.

Apply at the rate of 2 to 3 qts. per acre for **Pea Aphid** and **Pea Leaf Weevil**. For **Alfalfa Looper**, apply at the rate of 1 gal. per acre.

Use sufficient water to insure thorough coverage (5 to 10 gal. by air; 20 gal. by ground application). Repeat applications as needed to maintain control.

For the control of **Pea Weevil**, apply at the same rates as above Pea Aphid during early bloom period before eggs are laid. Repeat in 5 days.

Do not apply within 7 days of harvest.

Insist on ...

Miller

M and M S

INSECT

ALFALFA, CL

Active ingredients: Malathion
Methoxychlor, Technical
Methylated Naphthalene
Aromatic Petroleum Derivat
Inert ingredients: ...
O,O-dimethyl dithiophosphat
Equivalent to 20.416% 2,2,4-trichloroethane and 2,2,4-

CAUTION — Keep

See right side panel
Please read entire label. Use o



MILLER P
AGRICULTURAL PRODUC
PORTLAND.

Apply 2 qts per acre. Apply when eggs begin to hatch (in approximate application)

Peas and other legumes. Do not apply to alfalfa.

Pea Weevil,

alfalfa Looper,

and other insects (through coverage application) maintain control

Apply at the same time as the bloom period

Insist on...

Miller's



M and M SPRAY 22E

INSECTICIDE FOR

ALFALFA, CLOVER and PEAS

Active ingredients: Malathion (50% emulsifiable concentrate)	23.2%
Methoxychlor, Technical (95% emulsifiable concentrate)	23.2%
Methylated Naphthalene (90% emulsifiable concentrate)	2.6%
Aromatic Petroleum Derivative Solvents (90% emulsifiable concentrate)	45.0%
Inert ingredients	6.0%
O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate	
Equivalent to 20.416% 2,2-bis (p-methoxyphenyl)-1,1,1-trichloroethane and 2.754% ether isomers and reaction products	

CAUTION — Keep out of reach of children

See right side panel for additional cautions

Please read entire label. Use only as directed, noting all cautions.

1 GALLON

MILLER PRODUCTS CO.

AGRICULTURAL PRODUCTS DIVISION W. R. GRACE & CO
PORTLAND, OREGON, U. S. A.

CAUTION

Harmful if swallowed. Avoid prolonged breathing of spray. Avoid prolonged and repeated contact with skin. Wash thoroughly after using. Avoid contamination of feed and food.

Highly toxic to fish — do not contaminate lakes, ponds, or streams.

Do not apply when weather conditions favor drift from areas treated.

Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes or by burning (keep out of smoke).

NOTICE: Follow directions carefully. Timing and method of application may vary with crop conditions. Always use the minimum amount of spray. Do not apply to windward side of plants. Influencing factors in the use of this product are beyond the control of the manufacturer. We assume all risks of use, storage or handling of this material unless stated with directions on the container.

U.S. PATENT OFFICE REG. NO. 2,802,000