





This is a facsimile of a pre-printed bag
on which the attached label is applied.



Net weight 50 lbs.



CYPREX® - KARATHANE 42D

(4% Cyprex (Dodine) + 2% Karathane Dust)

AGRICULTURAL FUNGICIDE

Active ingredients

#2,4-Dinitro-6-octyl phenyl crotonate)	1.46%
#2,6-Dinitro-4-octyl phenyl crotonate).....	0.10%
#Nitrooctyl phenols (principally dinitro) ...	4.00%
Dodine (n-dodecylguanidine acetate)	94.44%
Inert ingredients	
#A mixture of 1-methylheptyl, 1-ethylhexyl, 1-propylpentyl isomers.	

E.P.A. Reg. No. 802-414-AA

*CYPREX is the Reg. T.M. of the American Cyanamid Co.

*KARATHANE is the Reg. T.M. of the Rohm & Haas Co.

CAUTION -- Keep out of reach of children.

Harmful if swallowed. Avoid breathing dust. May cause skin and eye irritation or sensitization. Avoid contact with skin and eyes. In case of contact with skin, wash thoroughly with soap and running water. For eyes, flush with plenty of water for at least 15 minutes and get medical attention.

Avoid contamination of feed or feedstuffs

This product is toxic to fish. Keep out of lakes, ponds and streams. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift of dust from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes.

Dispose of wastes by burying in non-crop lands away from water supplies. The containers should be disposed of by punching holes in them and burying with wastes.

APPLES, PEARS - Scab, Powdery Mildew

Apply 30 to 40 lbs. per acre at 5 to 10 day intervals or as needed to maintain scab and/or powdery mildew control from pre-bloss through petal fall and again in first cover application 10 days later. Use 20 lbs. in later cover applications as needed.

Do not apply within 21 days of harvest. Avoid drift of dust onto adjacent crops. Do not graze the cover crops in treated orchards.

USE PRECAUTIONS

Read entire label. Use strictly in accordance with label directions, cautions and warnings and in conformity with Federal and State regulations.

Treated apples and pears should not be used in the manufacture of animal feedstuffs or for use in livestock feeds. Such use may result in illegal residues in milk or meat.

Apply this product in accordance with this label. Do not apply when temperature will exceed 90 to 95° F. during day. Use extra care when to avoid drift onto plums, peaches, and bearing sweet cherries since severe foliage and fruit injury may result. The grower is responsible for residues on his crops as well as for damage caused by drift from his property to that of others.