DIRECTIONS FOR USE

Shake before using. Please read entire label. Use only as directed, noting all cautions. Consult State Agricultural Experiment Station or State Extension. Authorities for exact liming and spacing of sprays in your area.

WHERE TO USE	WHEN TO USE Tab	oiespoons pe al. of water
APPLES - Rust Mite (aids in control of)	Delayed Dormant	4
	Summer Use	3
APPLES, PEARS - San Jose Scale, Italian Pear Scale, Eggs of Aphids, Red Spider Mites.	Dormant or Delayed Dormant	5
Red Spider Mites	Summer Use	2
San Jose & Italian Pear Scales (Summer forms)	Summer Use	3
PEARS - Pear Psylla	Delayed Dormant	5
	Summer Use	3
Blister Mite	Delayed Dormant & Fall Spray	² 5
ARINES, PÉACHES, PLÜMS, PRUNES, WALNUTS, and ALMONDS. Scales: Brown Apricot, Black, San Jose, Olive (Parlatoria), Oleander, Frosted, Italian Pear European, Fruit Lecanium, Punam, Sott Brown: Eggs of Aphids, Red Spider Mites.	Dormant or Delayed Dormant	5
Red Spider Mites	Summer Use	5
Brown Apricot & San Jose Scales (Summer stages)	Summer Use	2 1/2
ORNAMENTALS AND SHADE TREES (Birch, Cherry (flowering), Dogwood, Evergreens (see Note),	Dermant Spray on Deciduous Varieties	5
Hawthorn, Holly, Magnolia, Maple, Oak, Peach (flowering), Privet, Pyracantha, Sycamores)	Dormant Spray on Evergreen Varieties	21/2
Red Spider Mites and Eggs, Scales: Soft Brown, Oyster Shell, Brown Apricot, Black and San Jose.	NOTE: May cause discoloration of Blue Spruce. Do not use on Evergreens or Ferns not tolera to oil sprays.	
FIGS - Fig Scale	Dormant or Delayed Dormant	7 1/2
OLIVES Spider Mites, Scales; Parlatoria, Oleander and Black	Prebloom or post bloom up to buckshot	• 21/2



DIRECTIONS FOR USE (continued)
WHERE TO USE WHEN TO USE

HERE TO USE WIT

Tablespoons per gal. of water

ORANGES (Valencia)
Red Spider Mites, Scale:
Red, Black, Yellow & Purple

Do not use later than Sept 1st.

it 1st. 21/2

CAUTION

⊮ ·Hanntut it swallowed! It swallowed do not induce vomiting. Call a physician immediately. Avoid inhalation of spray mist.

For pliage application the spray tank must be free of any sulphur residue the using this material. Do not combine sulphur sprays with this material place and do not follow a previous application of sulphur or apply sulphur following an application of this material until sufficient time has elapsed for safety (at least 30 days). If in doubt, spray a portion of one tree previously treated with sulphur, using 1 quart of oil to 25 gallons water. If sufficient time has not elapsed, burn will show in several days.

Do not use oil spray when temperatures are above 90°F. When temperatures drop below 32°F during night, do not spray tree and vine crops until daytime temperatures have warmed to a point above 35°F and crops are free from moisture due to melting frost or heavy dews.

Do not use on foliage closer then 60 days after or 90 days before using CAPTAN or PHALTAN* products. Do not use in combination with SEVIN*. Application to trees weakened by disease, drought, drying winds, or high nitrogen applications may result in oil injury.

ENVIRONMENTAL CAUTIONS

Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Do not reuse empty container. Wash thoroughly with water and detergent. Discard in safe place.

*PHALTAN is a registered trademark for phthalimide fungicide.
*SEVIN is the trademark of the Union Carbide Corp. for
1-naphthyl N-methylcarbamate.

NOTICE: Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

LC-7M

THE CHAS. H. LILLY CO. PORTLAND, OREGON 97218

EPA Est. 802-OR-1

EPA Reg. No 802-415