

LILLY/MILLER  
Superior Type  
SPRAY OIL

ACTIVE INGREDIENT:

Petroleum Oil . . . . 99%  
INERT INGREDIENTS . . . . 1%

Minimum Unsulfonated Residue 92%  
Petroleum Oils Classification - unclassified  
A 70 second viscosity type oil

Controls red spider mites  
For Dormant & Summer use

Keep out of reach of children  
CAUTION

See right side panel for additional cautions

NET CONTENTS 1 GALLON

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed, inhaled or absorbed through the skin.  
Avoid breathing spray mist. Avoid contact with skin, eyes or clothing.

STATEMENT OF PRACTICAL TREATMENT

If Swallowed: Do not induce vomiting. Call a physician or Poison  
Control Center immediately.

If Inhaled: Remove to fresh air.

If In Eyes: Flush with plenty of water. If irritation persists, get  
medical attention.

If On Skin: Wash with plenty of soap and water.

ENVIRONMENTAL HAZARDS

Do not apply directly to water. Do not apply when weather conditions  
favor drift from areas treated. Do not contaminate water by cleaning of  
equipment or disposal of wastes. Apply this product only as specified  
on this label.


DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner  
inconsistent with its labeling.

DIRECTIONS FOR USE

Shake well before using. Please read entire label. Use only as directed,  
noting all cautions. Consult State Agricultural Experiment Station or  
State Extension Authorities for exact timing and spacing of sprays in your  
area.

WHERE TO USE	WHEN TO USE	Tablespoons per gal. of water
APPLES - Rust Mite (aids in control of)	Delayed Dormant*	4
	Summer Use	3
APPLES, PEARS - San Jose Scale, Italian Pear Scale,	Dormant or Delayed Dormant*	5



Eggs of Aphids, Red Spider  
Mites

Red Spider Mites	Summer Use	2
San Jose & Italian Pear Scales (Summer forms)	Summer Use	3
PEARS - Pear Psylla	Delayed Dormant* Summer Use	5 3
Blister Mite	Delayed Dormant* & Fall Spray	5
APRICOTS, CHERRIES, NECTARINES, PEACHES, PLUMS, PRUNES, WALNUTS and ALMONDS - Scales: Brown Apricot, Black, San Jose, Olive (Parlatoria), Oleander, Frosted, Italian Pear, European Fruit Lecanium, Putnam, Soft Brown; Eggs of Aphids, Red Spider Mites	Dormant or Delayed Dormant*	5
Red Spider Mites	Summer Use	2
Brown Apricot & San Jose Scales (Summer stages)	Summer Use	2 1/2
ORNAMENTALS AND SHADE TREES Birch, Cherry (flowering), Dogwood, Evergreens (see note), Hawthorn, Holly, Magnolia, Maple, Oak, Peach (flowering), Privet, Pyracantha, Sycamores	Dormant or Delayed Dormant* Spray on Deciduous Varieties	5
Red Spider Mites and Eggs, Scales: Soft Brown, Oyster Shell, Brown Apricot, Black, and San Jose.	Dormant or Delayed Dormant* Spray on Evergreen Varieties	2 1/2
	NOTE: May cause dis- coloration of Blue Spruce. Do not use on Evergreens or Ferns not tolerant to oil sprays.	
FIGS - Fig Scale	Dormant or Delayed Dormant*	7 1/2
OLIVES - Spider Mites, Scales: Parlatoria, Oleander and Black	Pre-bloom or post bloom up to buckshot**	2 1/2
ORANGES (Valencia) Red Spider Mites, Scale: Red, Black, Yellow & Purple	Do not use later than Sept. 1st	2 1/2

- \*Delayed Dormant - The period just before blossom buds break open.
- \*\*Buckshot - Young olives about the size of peas.

NOTE: This product may be combined with Lilly/Miller Polysul Summer & Dormant Spray for use as a dormant combination spray. Refer to the Lilly/Miller Polysul label for recommended rates.

#### USE PRECAUTIONS

For foliage application the spray tank must be free of any sulphur residue before using this material. Do not combine sulphur sprays with this material on foliage during spring and summer months and do not follow a previous application of sulphur or apply sulphur following an application of this material until sufficient time has elapsed for safety (at least 30 days). If in doubt, spray a portion of one tree previously treated with sulphur using 1 quart of oil to 25 gallons of water. If sufficient time has not elapsed, burn will show in several days.

Do not use oil spray when temperatures are above 90 degrees F. When temperatures drop below 32 degrees F during night, do not spray tree and vine crops until daytime temperatures have warmed to a point above 35 degrees F and crops are free from moisture due to melting frost or heavy dews.

Do not use on foliage closer than 60 days after or 90 days before using CAPTAN or PHALTAN\* products. Do not use in combination with SEVIN\*. Application to trees weakened by disease, drought, drying winds, or high nitrogen applications may result in oil injury.

\*PHALTAN is the registered trademark for phthalimide fungicide.  
\*SEVIN is the reg. T.M. of the Union Carbide Corporation for carbaryl insecticide.

#### STORAGE AND DISPOSAL

Store this product in its original container and keep in a locked storage area out of reach of children and domestic animals.

Do not reuse container. Rinse thoroughly before discarding and securely wrap original container in several layers of newspaper and discard in trash.

NOTICE: Follow directions carefully. Timing and method of application, weather or crop conditions, mixtures with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

THE CHAS. H. LILLY CO.  
PORTLAND, OREGON 97218  
EPA Est. 802-OR-1  
EPA Reg. No. 802-415  
79756