

ACCEPTED
 Oct 6, 1969
 U.S. DEPARTMENT OF AGRICULTURE
 OFFICE OF PLANT INDUSTRY
 WASHINGTON, D.C. 20250

KELTHANE-DIAZINON* 42D

(4% Kelthane* + 2% Diazinon* Dust)

AGRICULTURAL INSECTICIDE

Active ingredients	
1,1-bis (chlorophenyl)-2,2,2-trichloroethanol. . .	4%
0,0-diethyl O-(2-isopropyl-4-methyl-6-pyrimidinyl) phosphorothioate†.	2%
Inert ingredients	94%
† U.S. Pat. No. 2,754,243	

CAUTION— Keep out of reach of children.

HARMFUL IF SWALLOWED! May be absorbed through skin. Avoid inhalation of dust. Avoid contact with skin, eyes or clothing. Wash thoroughly after handling.

Avoid contamination of feed and foodstuffs. Do not apply when weather conditions favor drift from areas treated.

Apply this product only as specified on this label.

This product is toxic to fish and wildlife. Keep out of lakes, streams and ponds.

Birds feeding on treated area may be killed.

Dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes, or by burning (stay out of smoke and fumes).

GENERAL DIRECTIONS

Apply the specified dosage for the specific crop as listed. Make first application when insects first appear and repeat as necessary for adequate control. Observe use precautions.

*Kelthane is the Reg. T.M. of the Rohm & Haas Co.
 †Diazinon is the Reg. T.M. of Geigy Agricultural Chemicals.

SPECIFIC CROPS

BEANS (Snap, Pole, Lima) - Aphids, Cucumber Beetle, Serpentine Leaf Miners, Two-spotted and Pacific Mites
 USE 30 lbs. per acre. Do not apply within 7 days of harvest. Do not feed treated vines to poultry, dairy or meat animals.

CANTALOUPE, CUCUMBERS, WINTER & SUMMER SQUASH, WATERMELONS
 Aphids, Serpentine Leaf Miners, Thrips, Two-spotted and Pacific Mites
 USE 30 lbs. per acre. Do not apply to Summer Squash and Cucumbers within 7 days of harvest; to the others listed within 3 days of harvest.

TOMATOES, PEPPERS - Aphids, Serpentine Leaf Miners, Pacific, Two-spotted and Tomato Russet Mites
 USE 30 to 35 lbs. per acre. Do not apply to Tomatoes within 2 days of harvest; to Peppers within 5 days.

HOPS - Two-Spotted Mites
 USE 25 to 35 lbs. per acre. Do not apply within 14 days of harvest.

STRAWBERRIES - Aphids, Atlantic Mites
 USE 30 to 35 lbs. per acre. Do not apply within 5 days of harvest.

NOTICE: Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other pesticides, and other factors in the use of this product are beyond the control of the manufacturer. Risks of use, storage or handling of the material are not according with directions given herewith.