

KELTHANE*-SULPHUR 450D

(4% Kelthane* - 50% Sulphur Dust)

AGRICULTURAL INSECTICIDE

Active ingredients	
1,1-bis (chlorophenyl)-2,2,2-trichloroethanol	4.0%
Sulphur	50.0%
Inert ingredients	46.0%

APPLICATION DIRECTIONS: Apply on first appearance of mites and repeat if necessary for adequate control.
 †Do not apply closer to harvest than number of days indicated for the specific crop being treated.

CAUTION— Keep out of reach of children.

Avoid skin contact. Avoid inhalation of dust. Wash thoroughly after using.

Avoid contamination of feed and foodstuffs.

USE PRECAUTIONS

Read entire label. Use strictly in accordance with label directions, cautions, and warnings and in conformity with Federal and State regulations.

Avoid drift to adjoining food and forage crops about to be harvested. Apply this product only as specified on this label. Do not apply when weather conditions favor drift of dust from areas treated. The grower is responsible for residues on his crops as well as for damages caused by drift from his property to that of others.

Do not apply when shade temperatures exceed or are likely to exceed 90° F. as sulphur may burn foliage.

Dispose of wastes by burying in non-crop lands away from water supplies. The containers should be disposed of by punching holes in them and burying with wastes. Do not contaminate water by cleaning of equipment, or disposal of wastes.

This product is toxic to fish. Keep out of lakes, streams and ponds. Do not apply where runoff is likely to occur.

Crop and Pest	Pounds	†Days per Acre
DALFALFA - Seed Crops Only		
Two-spotted and Pacific Mites	0	25
Use restricted to Grant & Adams Counties in Washington and to California. Fields not to be grazed by livestock. Crop cuttings or residues not to be fed or baled. The fields must be cleared after harvest by burning or by discing into the soil.		
BEANS - Dry, Green and Lima		
Two-spotted and Pacific Mites	7	25 to 30
Do not feed treated vines to meat or dairy animals.		
HOPS - Two Spotted Mites	7	25 to 35
TOMATOES, EGGPLANTS, PEPPERS - Two-spotted, Pacific and Tomato Russet Mites		
	2	25 to 35

NOTE: If crop to be treated will be used for canning or freezing purposes, consult food processor before applying sulphur.

ACCEPTED
 MAY 9 1973
 UNDER THE...
 FEDERAL...
 REGULATIONS

NOTICE: Follow directions carefully. Timing and method of application, weather, and crop conditions, mixtures with other pesticides, and other factors in the use of this product are beyond the control of the manufacturer. Buyer assumes all risks of use, storage or handling of this material. For full assurance with directions given hereon.