

 **LILLY/MILLER** BEST DOCUMENT AVAILABLE

# 10% DITHANE\* M-45 DUST

**ACTIVE INGREDIENTS:**

Zinc ion and manganese ethylene bisdithiocarbamate 80%, a coordination product of manganese 16%, zinc 2%, ethylene bisdithiocarbamate 62% .....

INERT INGREDIENTS ..... 92%  
Total ..... 100%

Keep out of reach of children

## CAUTION

### PRECAUTIONARY STATEMENTS

#### HAZARD TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** May cause irritation of nose, throat, eyes and skin. Do not breathe dust. Protective clothing such as long sleeve shirt, long pants, rubber gloves and rubber boots are to be worn by persons involved in mixing and loading of this product

In case of skin contact, wash with water; for eyes, wash copiously with water and get medical attention.

#### ENVIRONMENTAL HAZARD

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

### DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

**NET WT. 50 LBS.**

THE CHAS. H. LILLY CO.  
Portland, Oregon 97210  
EPA Reg. No. 802-428  
EPA Est. 802-OR-1

 **LILLY/MILLER**

### STORAGE AND DISPOSAL

Keep away from fire and sparks. Store in a cool dry place.

Keep container closed when not in use. Do not contaminate water, food or feed by storage or disposal.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

### DIRECTIONS FOR USE

#### POTATO SEED PIECE TREATMENT

for protection against *Fusarium sambucinum*.

Lilly/Miller 10% DITHANE M-45 DUST is a special fungicide formulation for application on whole or cut potato tubers to protect them from seed-piece decay caused by *Fusarium sambucinum*.

Use 1 lb. of dust per 100 lbs. of whole or cut potato seed pieces. Apply the dust in a manner such that the entire surface of the seed-piece becomes thoroughly covered with the fungicide dust. Plant seed as soon after treatment as possible.

**DO NOT USE TREATED SEED PIECES FOR FOOD OR FEED PURPOSES.**

\*DITHANE is the Reg. T.M. of the Rohm & Haas Co.

**NOTICE:** Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material, not in strict accordance with directions given herewith.

LC-2S