Miller's ROSE AND FLOWER SPRAY is a highly effective combination Insecticide, Miticide and Fungicide for Roses, Flowers and Ornamentals (Azaleas, Asters, Carnations, Chrysanthemums, Delphiniums, Fuchsias, Geraniums, Gladiolus, Marigolds, Snapdragons, Zinnias).

Please read entire label. Use only as directed, noting all Cautions and Warnings.

DIRECTIONS FOR USE

To operate


- 1. Shake container before using.
- 2. **IMPORTANT:** Hold can upright and **18 to** 24 inches from plant. Apply as a light spray. Only a fine mist is required.
- 3. Spray with slow, sweeping motion, covering about one foot of plant per second, or spray short spurts at intervals. Be sure spray contacts both upper and lower sides of leaves. Spray preferably in early morning or evening and, if possible, when the air is still.
- 4. Avoid spraying in direct sunlight, during heat of day, or when plants are wet. Examine plants often and spray at first sign of infestation.

PLANT DISEASES

To prevent Powdery Mildew and Black Spot cf Roses and certain Leaf Spots of other flowers (such as Septoria Leaf Spot of Chrysanthemums and Carnations), spray as directed above at weekly intervals. When heavy infection occurs, spray every 4th day. For best results apply at weekly intervals before disease appears.

PLANT INSECTS --- Aphids, Thrips (exposed), Spider Mites, Lace Bugs, Leafhoppers, Soft Brown Scale (crawler stage), Mealy Bugs, White Fly

On first sign of insect infestation, spray as directed above at weekly intervals. When heavy infestations occur, spray twice weekly. For Soft Brown Scale (crawler stage), Mealy Bugs, White Fly, make repeated applications, taking special care to contact insects with spray bursts of about one secund (time).





· .

CAUTION Keep out of the reach of children See right side panel for additional cautions E.P.A. Reg. No. 802-433-A

CAUTION

Harniful if swallowed, inhaled or absorbed through skin. Avoid inhalation of spray mist Avoid contact with skin and eyes Wash thoroughly after using. May be absorbed through skin. In case of contact, wash immediately with soap and water.

Avoid contamination of feed and foodstuffs This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Do not use in confined area. Do not apply during heat of day or in direct sunlight. Do not apply to wet plants. Do not use on Maidenhair Fern, Lantana or Cyclamen; or on Vegetables.

ENVIRONMENTAL CAUTIONS

This product is toxic to fish, birds, and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams, or ponds, Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes Apply this product only as specified on this label.

WARNING – Contents under Pressure

Do not puncture or incinerate container. Never throw container into fire

Do not expose container to heat or store at temperatues above 120° F. Keep at room terni perature - away from direct surilight, radiators, stoves, hot water and other heat

A TURN AF FNTS	
(a)	· · · · · · · · · · · · · · · · · · ·
ter an fair an the training the training the state of the	r 225°•
(a) A set of the s	ta ng kana
	(, ≭)(0, 0 (), 2()t (
the state the second	
n an geologica d'Altante. An geologica de la companya de la co An antica de la companya de la compa	÷ • •
	, <i>1 , j ; , ,</i> ,

A second sec second sec

**Piper in the second to are the Piperpiper to a

NOTICE. For A day to a construct the property of the and the second والمراجع والمراجع والمواجع والمتعام والمراجع والمراجع والمراجع والمراجع والمراجع in the construction of the group, that because prove out the (a) a set of a se

THE CHAS HELLEY CO. and the second TELLER SOLOR 1