

*Miller's*  
**CHLORDANE**  
**10% DUST**

PLEASE READ ENTIRE LABEL. USE ONLY AS DIRECTED,  
NOTING ALL CAUTIONS AND WARNINGS.

**DIRECTIONS**

**THRIPS, Certain CATERPILLARS, SOWBUGS, ANTS, LEAF MINERS, LEAFROLLERS, and certain other insects on FLOWERS**

1. Dust plants lightly, but thoroughly, being sure to contact both sides of the leaves.

**CABBAGE LOOPERS, FLEA BEETLES, SOWBUGS, COLORADO POTATO BEETLES, TOMATO HORNWORMS, THRIPS, LEAF MINERS and certain other garden insects on BEANS, CARROTS, CABBAGE, CUCUMBERS, POTATOES, TOMATOES**

1. Dust lightly being sure to contact both top and lower leaf surfaces.
2. Note . . . Do not apply after edible portions have begun to form. Do not apply to leafy vegetables such as Lettuce or Spinach. See Caution.

**ANTS**

1. As an area treatment, use 1 lb. to 400 sq. ft.
2. Dust in and along cracks and crevices in foundations and walls, around nests and hills, in runways, and around foundations and outbuildings.
3. Spot treat soil or lawns in heavily infested areas.
4. Do not use in homes, in dairy buildings, in barns, or poultry houses.

**EARWIGS**

1. Use 1 lb. to 1000 sq. ft.
2. Apply to ground around edges of lawns, buildings and foundations, around and under rocks, boards and other damp shady places where Earwigs are found.

**GRASSHOPPERS, CRICKETS, EARWIGS, ANTS on Tomatoes, Cabbage, Broccoli, Potatoes, Gladioli, Zinnias**

1. Dust plants and surrounding ground lightly.
2. Note . . . Do not apply after edible portions have begun to form.

**SOD WEBWORM (Lawn Moth)**

1. Dust lawn using 1 lb. to 500 to 800 sq. ft.
2. Sprinkle to wash into soil after applying.
3. After application has been made, do not permit children and pets on treated area until the insecticide has been washed into the soil by sprinkling and the grass has completely dried. Birds feeding on treated areas may be harmed.

**CAUTION** . . . Avoid inhalation. Wash hands after use. Keep out of reach of children. Do not contaminate drinking water or use near fishponds, etc.


Avoid contamination of feed and foodstuffs. Remove residues remaining at harvest by washing, wiping, or other effective means. Do not apply to cucurbits that may be susceptible to injury by chlordane. Do not apply to green or spring onions; to leafy vegetables; nor to tomatoes, cucumbers or squash after fruit begins to form. Do not . . .

NET WT.  
2 LBS.

\$1.75

  
*Miller's*

**Chlordane**  
**10% Dust**



1. As an area treatment, use 1 lb. to 400 sq. ft.
2. Dust in and along cracks and crevices in foundations and walls, around nests and hills, in runways, and around foundations and outbuildings.
3. Spot treat soil or lawns in heavily infested areas.
4. Do not use in homes, in dairy buildings, in barns, or poultry houses.

**EARWIGS**

1. Use 1 lb. to 1000 sq. ft.
2. Apply to ground around edges of lawns, buildings and foundations, around and under rocks, boards and other damp shady places where Earwigs are found.

**GRASSHOPPERS, CRICKETS, EARWIGS, ANTS on Tomatoes, Cabbage, Broccoli, Potatoes, Gladioli, Zinnias**

1. Dust plants and surrounding ground lightly.
2. Note . . . Do not apply after edible portions have begun to form.

**SOD WEBWORM (Lawn Moth)**

1. Dust lawn using 1 lb. to 500 to 800 sq. ft.
2. Sprinkle to wash into soil after applying.
3. After application has been made, do not permit children and pets on treated area until the insecticide has been washed into the soil by sprinkling and the grass has completely dried. Birds feeding on treated areas may be harmed.

**CAUTION** . . . Avoid inhalation. Wash hands after use. Keep out of reach of children. Do not contaminate drinking water or use near fishponds, etc.

Avoid contamination of feed and foodstuffs. Remove residues remaining at harvest by washing, wiping, or other effective means. Do not apply to cucurbits that may be susceptible to injury by chlordane. Do not apply to green or spring onions; to leafy vegetables; nor to tomatoes, cucumbers or squash after fruit begins to form. Do not use on Cole Crops after heads or other edible parts begin to form; nor on green or string beans after pods begin to form.

Do not apply on plants in bloom, as chlordane is toxic to pollenizing insects. Do not feed forage treated with this material to dairy animals or to animals being finished for slaughter.

Do not apply to lawn or other turf grasses when people or animals are on them and do not allow dust to drift to areas where it might injure people or animals or contaminate feed or food.

Toxic to fish and wildlife. Keep out of any body of water. Birds feeding on treated areas may be harmed. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.


Destroy empty container. Do not re-use for any purpose.

**NOTICE:** Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risk of use, storage or handling of this material not in strict accordance with directions given herewith.

MP 12F

**MILLER PRODUCTS**  
Portland, Oregon 97218

# 10% Dust



**Active Ingredients**  
Chlordane, technical\* . . . . . 10%  
**Inert Ingredients** . . . . . 90%

\*Equivalent to 6% octachloro-4,7 methano-tetrahydroindane and 4% related compounds.

**CAUTION** Keep out of the reach of children  
See back panel for additional cautions.  
E.P.A. Reg. No. 802-436-AA